arXiv:1207.4844v1 [math.CA] 20 Jul 2012

Exact Hausdorff and packing measures of linear Cantor sets with overlaps

QIU Hua

Department of Mathematics, Nanjing University, Nanjing, 210093, China

Email: huatony@gmail.com

Abstract.

Let K be the attractor of a linear iterated function system (IFS) $S_j(x) = \rho_j x + b_j$, $j = 1, \cdots, m$, on the real line satisfying the generalized finite type condition, (whose invariant open set $\mathcal{O}$ is an open interval) with an irreducible weighted incidence matrix. This condition is introduced by Lau & Ngai recently as a natural generalization of the open set condition, allowing us to include many important overlapping cases. It is known that the Hausdorff and packing dimensions of K coincide and can be calculated in terms of the spectral radius of the weighted incidence matrix. Let α be the dimension of K. In this paper, we state that

$$\mathcal{H}^\alpha(K \cap J) \le |J|^\alpha$$

for all intervals $J \subset \overline{\mathcal{O}}$, and

$$\mathcal{P}^\alpha(K \cap J) \ge |J|^\alpha$$

for all intervals $J \subset \overline{\mathcal{O}}$ centered in K, where $\mathcal{H}^\alpha$ denotes the α-dimensional Hausdorff measure and $\mathcal{P}^\alpha$ denotes the α-dimensional packing measure. This result extends a recent work of Olsen in which the open set condition is required. We use these inequalities to obtain some precise density theorems for the Hausdorff and packing measures of K. Moreover, using these densities theorems, we describe a scheme for computing $\mathcal{H}^\alpha(K)$ exactly as the minimum of a finite set of elementary functions of the parameters of the IFS. We also obtain an exact algorithm for computing $\mathcal{P}^\alpha(K)$ as the maximum of another finite set of elementary functions of the parameters of the IFS. These results extend previous ones by Ayer & Strichartz and Feng, respectively, and apply to some new classes allowing us to include linear Cantor sets with overlaps.

1 Introduction and Statement of Results

In this paper we will analyze the bahaviour of the Hausdorff and packing measures of self-similar sets satisfying the generalized finite type condition which is weaker than the

---

MR(2000) Subject Classification: 28A78, 28A80

Supported by NSFC 10901081

open set condition. In particular, we will deal with the exact calculating of the Hausdorff and packing measures for a special kind of linear Cantor sets.

The problem of calculating the Hausdorff and packing dimensions of the attractor of an self-similar iterated function system (IFS) is one of the most interesting questions in fractal geometry. During the past two decades there has been an enormous body of literatures investigating the Hausdorff and packing dimensions of self-similar sets and wide ranging generalizations thereof. See the books [2], [4], [13] and the references therein. Let $\{S_j\}_{j=1}^m$ be an IFS of contractive similitudes on $\mathbb{R}^d$ defined as

$$S_j(x) = \rho_j R_j x + b_j, \quad j = 1, \cdots, m, \tag{1.1}$$

where for all j, $0 < \rho_j < 1$ is the contraction ratio, R_j is an orthogonal transformation and $b_j \in \mathbb{R}^d$. Let K denote the *self-similar set* (or *attractor*) of the IFS, i.e., K is the unique non-empty compact set in $\mathbb{R}^d$ satisfying

$$K = \bigcup_{j=1}^{m} S_j(K).$$

A basic result is that the Hausdorff dimension $\dim_H K$ and the packing dimension $\dim_P K$ are always equal for the self-similar set K, i.e.,

$$\dim_H K = \dim_P K,$$

(see [3]). In general it is quite difficult to calculate the common value of the Hausdorff and packing dimensions of the attractor K. It is a classical result (see Moran [16], Hutchinson [7]) that if the IFS satisfies the *open set condition*, i.e., there exists a non-empty bounded open set $\mathcal{O} \subset \mathbb{R}^d$ such that $\bigcup_{j=1}^m S_j(\mathcal{O}) \subset \mathcal{O}$ and $S_i(\mathcal{O}) \cap S_j(\mathcal{O}) = \emptyset$ for all $i \neq j$, the common value of the Hausdorff dimension $\dim_H K$ and the packing dimension $\dim_P K$ of K is the unique solution α of the equation

$$\sum_{j=1}^{m} \rho_j^{\alpha} = 1. \tag{1.2}$$

Non-overlapping or almost non-overlapping self-similar IFSs have been studied in great detail via the open set condition.

In the absence of the open set condition, much less is known about IFSs with overlaps. To deal with such systems, by extending a method of Lalley [8] and Rao &Wen [21], Ngai & Wang [18] formulated a weaker separation condition, i.e., the *finite type condition*, which may includes many important overlapping cases and described a method for computing the Hausdorff and packing dimensions of the attractor in terms of the spectral radius of a certain weighted incidence matrix. The finite type condition requires the contraction ratios of the IFS's maps to be exponentially commensurable and thus does not generalize the open set condition. Recently, Lau and Ngai [9] introduced a more general condition, the *generalized finite type condition*, that extends both the open set condition and the finite type condition. Moreover, it does not require the contraction similitudes in the IFS to have exponentially commensurable contraction ratios as in the original finite type condition. Under such a generalized finite type condition, one can also compute the Hausdorff and packing dimensions of the attractor in terms of the spectral radius of a certain weighted incidence matrix.

Another central problem concerning the theory of self-similar IFSs is to estimate the Hausdorff and packing measures of self-similar sets, which is also an area of active research [1, 6, 11, 12, 25]. In these papers and the references therein one can find the analysis of the estimation of the values of the Hausdorff or packing measures of some particular self-similar constructions. Since the definitions of the Hausdorff and packing measures are sometimes awkward to work with, there are only very few non-trivial examples of sets whose exact Hausdorff measures or packing measures are known. [25] is a recent review of relevant open questions in this field.

So far all these researches have been mainly addressed to the determination of the upper and lower bounds of the Hausdorff and packing measures. With regard to the determination of the exact values of the Hausdorff or packing measures, two papers [1] and [6] should be mentioned.

In [1], Ayer and Strichartz considered a kind of linear Cantor set K which is the attractor of a linear IFS $S_j(x) = \rho_j x + b_j$, $j = 1, \cdots, m$, on the real line satisfying the open set condition (where the open set is the open interval (0,1)). They gave an algorithm for computing the α-dimensional Hausdorff measure $\mathcal{H}^\alpha(K)$ exactly as the minimum of a finite set of elementary functions of the parameters of the IFS by using the fact that the exact α-dimensional Hausdorff measure $\mathcal{H}^\alpha(K)$ is the inverse of the maximal density of intervals contained in $[0, 1]$ with respect to the α-dimensional normalized Hausdorff measure λ of $\mathcal{H}^\alpha$ restricted to K, i.e., $\lambda = \mathcal{H}^\alpha|_K/\mathcal{H}^\alpha(K)$, where α is the common value of the Hausdorff and packing dimensions of K. It should be pointed out that if the open set condition is satisfied, the α-dimensional Hausdorff measure restricted to K and the α-dimensional packing measure restricted to K are proportional. Hence the α-dimensional normalized Hausdorff measure λ is also equal to $\mathcal{P}^\alpha|_K/\mathcal{P}^\alpha(K)$ where $\mathcal{P}^\alpha$ denotes the α-dimensional packing measure.

On the other hand, in [6], Feng proved that the α-dimensional packing measure $\mathcal{P}^\alpha(K)$ of K is equal to the inverse of the so-called minimal centered density of intervals centered in K with respect to the α-dimensional normalized Hausdorff measure λ, which also yields an explicit formula for calculating the exact α-dimensional packing measure $\mathcal{P}^\alpha(K)$ of K in terms of the parameters of the IFS.

However, in these papers, one needs to work on self-similar sets with open set condition, which is a canonical class of fractals that plays an important role in the development of fractal geometry. To the best of our knowledge, there is no result concerning the exact Hausdorff or packing measures of self-similar sets without open set condition. Since the calculation of the Hausdorff and packing dimensions of self-similar sets with the open set condition can be successfully extended to those sets satisfying the generalized finite condition which includes many interesting overlapping cases, and in view of the above discussion, it is natural to ask whether the Hausdorff or packing measure of the linear Cantor set K with only the generalized finite type condition can also be calculated exactly. This is the motivation of this paper.

Recall that in order to get the exact values of the Hausdorff and packing measures of the linear Cantor set K with open set condition (where the open set is the open interval (0,1)). The following explicit formulae for the Hausdorff measure $\mathcal{H}^\alpha(K)$ and the packing measure $\mathcal{P}^\alpha(K)$ of K play a key role.

$$\mathcal{H}^\alpha(K)^{-1} = \sup\{\frac{\lambda(J)}{|J|^\alpha} : J \text{ is an interval with } J \subset [0,1]\}, \tag{1.3}$$

and

$$\mathcal{P}^\alpha(K)^{-1} = \inf\{\frac{\lambda(J)}{|J|^\alpha} : J \text{ is an interval centered in } K \text{ with } J \subset [0,1]\}, \tag{1.4}$$

where α is the common value of the Hausdorff and packing dimensions and λ denotes the α-dimensional normalized Hausdorff measure restricted to K. Formula (1.3) was implicit in earlier work by Marion [11, 12] and in [1] by Ayer & Strichartz, while formula (1.4) was proved in a direct and elementary way in [6] by Feng.

Recently, Morán [17] and Olsen [19] generalized the above two formulae to general case independently. In [17] the so-called self-similar tiling principle plays a central role in the proof. This principle says that any open subset U of K can be tiled by a countable set of similar copies of an arbitrarily given closed set with positive Hausdorff or packing measure while the tiling is exact in the sense that the part of U which cannot be covered by the tiles is of null measure.

However, the proof in [19] is quite different from that in [17]. Let K be a self-similar set in $\mathbb{R}^d$ as described in (1.1) with the Hausdorff and packing dimensions of K equal to α. Consider K which satisfies the open set condition or the strong separation condition. Recall that in [19] Olsen perform a detailed analysis of the behaviour of the Hausdorff measure $\mathcal{H}^\alpha(K \cap U)$ and of the packing measure $\mathcal{P}^\alpha(K \cap B(x, r))$ of small convex Borel sets U and balls $B(x, r)$. In particular, he showed that if K is a self-similar set satisfying the open set condition,

$$\mathcal{H}^\alpha(K \cap U) \le |U|^\alpha$$

for all convex Borel sets U. A dual result for the α-dimensional packing measure was also proved which says that

$$\mathcal{P}^\alpha(K \cap B(x, r)) \ge (2r)^\alpha$$

for all $x \in K$ and all small $r > 0$ if K satisfies the strong separation condition. The latter result was also generalized to the open set condition by the author recently (see [20]) in order to prove the continuity of packing measure function of self-similar IFSs. In fact, the above inequality concerning the α-dimensional packing measure also holds for each $B(x, r)$ contained in $\mathcal{O}$ with $x \in K$, where $\mathcal{O}$ is the open set required in the open set condition with $\mathcal{O} \cap K \neq \emptyset$. (There must exist such $\mathcal{O}$ since the open set condition is equivalent to the strong open set condition. See [22].)

To make the ideas clear and to match our question precisely we restrict our interest to the linear Cantor set K defined before. Hence the above two formulae are rewroted in the following form, i.e.,

$$\mathcal{H}^\alpha(K \cap J) \le |J|^\alpha \tag{1.5}$$

for all intervals $J \subset [0, 1]$, and

$$\mathcal{P}^\alpha(K \cap J) \ge |J|^\alpha \tag{1.6}$$

for all intervals $J \subset [0, 1]$ centered in K where α is the common value of the Hausdorff dimension and the packing dimension.

As an application of (1.5) and (1.6), Olsen also proved formulae (1.3) and (1.4) (Actually, he proved the general versions of them.) through the classical density theorems of geometric measure theory which were stated for arbitrary subsets of Euclidean space [2, 13]. Formulae (1.3) and (1.4) show that the exact values of the Hausdorff and packing measures of K coincide with the inverses of the supremum and infimum of the densities of the α-dimensional normalized Hausdorff measure λ on suitable classes of sets, respectively.

In order to calculate the exact values of the Hausdorff and packing measures of the linear Cantor set K satisfying only the generalized finite type condition, we need to establish similar explicit formulae as (1.3) and (1.4). Following the frame of Olsen's work, two inequalities similar to (1.5) and (1.6) are needed. Recall that in proving (1.5) and (1.6), one should find optimal coverings and packings in a self-similar setting which require almost non-overlap among the various similar pieces into which the fractal decomposes. In view of this, it is therefore entirely plausible that the open set condition is indispensible. In the present paper, somewhat surprisingly, we will show that the formulae (1.5) and

(1.6) still hold under the assumption that the weighted incidence matrix of K is irreducible where K is required to satisfy only the generalized finite type condition. This leads to the following results.

Theorem 1.1. *Let K be a linear Cantor set satisfying the generalized finite type condition with respect to the invariant open set $(0,1)$ with an irreducible weighted incidence matrix A_α, where α is the Hausdorff dimension of K. Then*

$$\mathcal{H}^\alpha(K \cap J) \le |J|^\alpha \tag{1.7}$$

for all intervals $J \subset [0,1]$.

Theorem 1.2. *Let K be the linear Cantor set described as before. Then*

$$\mathcal{P}^\alpha(K \cap J) \ge |J|^\alpha \tag{1.8}$$

for all intervals $J \subset [0,1]$ centered in K.

We will give the detailed definition of the generalized finite type condition and the exact concept of the weighted incidence matrix A_α in Section 2.

The idea of establishing Theorem 1.1 and Theorem 1.2 is the following. We first observe that if the weighted incidence matrix A_α of K is irreducible, K can be decomposed into an union of a set K_a with a graph directed construction and an attractor K_b of a countable infinite IFS with the open set condition (see [14], [15] for further properties of the graph directed sets and the infinite IFSs, respectively). Moreover, the common value of the Hausdorff and packing dimensions of K_a is strictly less than that of K_b. Hence the subset K_a will have null α-dimensional Hausdorff measure and null α-dimensional packing measure which ensures us to consider K_b in stead of K. Here α is the common value of the Hausdorff and packing dimensions of K (and also K_b, of course). Noticing that K_b is an attractor of a countable infinite IFS satisfying the open set condition, it is possible to adapt the techniques for proving (1.5) and (1.6) to establish Theorem 1.1 and Theorem 1.2.

By a similar discussion as that for classical self-similar sets with open set condition, the α-dimensional Hausdorff measure restricted to K_b and the α-dimensional packing measure restricted to K_b are also proportional. Obviously the above fact still holds if we replace K_b by K. We still write the α-dimensional normalized Hausdorff measure restricted to K as λ, then $\lambda = \mathcal{H}^\alpha|_K/\mathcal{H}^\alpha(K) = \mathcal{P}^\alpha|_K/\mathcal{P}^\alpha(K)$. We will show the λ-measure of some special kind of sets called islands of K can be expressed in terms of the parameters of the IFS of K. Then following the frame of Olsen's work, we use the inequalities (1.7) and (1.8) to get the explicit formulae of the Hausdorff and packing measures. In particular, we show that the exact value of these measures coincides with the inverse of the supremum or the infimum of the densities of λ on suitable classes of sets, respectively.

Corollary 1.3. *Let K be the linear Cantor set described as before. Then*

$$\mathcal{H}^\alpha(K)^{-1} = \sup\{\frac{\lambda(J)}{|J|^\alpha} : J \text{ is an interval with } J \subset [0,1]\}. \tag{1.9}$$

Corollary 1.4. *Let K be the linear Cantor set described as before. Then*

$$\mathcal{P}^\alpha(K)^{-1} = \inf\{\frac{\lambda(J)}{|J|^\alpha} : J \text{ is an interval centered in } K \text{ with } J \subset [0,1]\}. \tag{1.10}$$

These corollaries extend the results in [17, 19, 20]. Following the technique frame of [1, 6], under suitable assumptions, we then give an algorithm in theory for computing the α-dimensional Hausdorff measure $\mathcal{H}^\alpha(K)$ and the α-dimensional packing measure $\mathcal{P}^\alpha(K)$ exactly as the inverse of the maximal or minimal value of suitable finite sets of elementary functions of the parameters of the IFS respectively. This is possible since we could make a detailed analysis of λ, and thus a detailed analysis of the supremum in (1.9) and the infimum in (1.10) respectively. It should mention that we may allow touching islands, and indeed this case will lead to some of complicated and interesting phenomena. Due to the fact that the self-similar construction of K with the generalized finite type condition is much more complicated than that with the open set condition, our description of the exact calculations of the two kinds of measures will need some new important notations and techniques. We will describe a big scheme for the exact computing, which is a major adaption of the techniques frame used in [1] and [6].

This paper is organized as follows. In Section 2, we give some notations and basic facts about the generalized finite type condition. Our description of the generalized finite type condition is slightly different but equivalent to the original version in [9]. In Section 3, we deal with the density theorems for the Hausdorff and packing measures of linear Cantor sets satisfying the generalized finite type condition. Firstly, we give the proofs of Theorem 1.1 and Theorem 1.2 respectively, which will play an important role in giving the explicit formulae of the Hausdorff and packing measures. Secondly, we prove the formulae in Corollary 1.3 and Corollary 1.4 using the classical density theorems of geometry measure theory. Throughout this section and the following ones, we will always assume that the weighted incidence matrix of K is irreducible. In Section 4, we focus on the calculation of the exact Hausdorff measure and packing measure of the linear Cantor K under some suitable assumptions. A big scheme is provided. Section 5 collects some further discussions on this subject. We consider the possibility of dropping some assumption required in Section 4. We discuss briefly the slightly more general cases of IFSs that contain orientation reversing simlarities. We also consider the situation in higher dimensional Euclidean spaces and show why our results can not be generalized. Throughout the context, we will show some interesting and non-trivial examples.

2 Preliminaries

For convenience, we introduce a slightly different but equivalent description of the original definition of the generalized finite type condition in [9]. We focus the interest on the linear Cantor sets on the real line $\mathbb{R}$.

We will use the following notations throughout the paper. For a Borel measure ν on $\mathbb{R}$ and a Borel set E, we let $\nu|_E$ denote the restriction of ν to E, i.e.,

$$\nu|_E(U) = \nu(E \cap U)$$

for all Borel sets U. For any subset $E \subset \mathbb{R}$, we denote the *diameter* of E by $|E|$. For any $x \in \mathbb{R}$, let $\operatorname{dist}(x, E)$ denote the distance between x and E. That is

$$\operatorname{dist}(x, E) = \inf\{|x - y| : y \in E\}.$$

If A is any finite or countable set, we denote by $\sharp A$ the *cardinality* of A. We need the definitions of Hausdorff measure and Hausdorff dimension, packing measure and packing dimension. For $E \subset \mathbb{R}^d$, $s \geq 0$ and $\delta > 0$, put $\mathcal{H}^s_\delta(E) := \inf\{\sum_i |U_i|^s\}$, where the infimum is taken over all *δ-coverings* of E, i.e., countable collections $\{U_i\}$ of subsets of $\mathbb{R}^d$ with diameters smaller than δ and such that $E \subset \bigcup_i U_i$. The *s-dimensional Hausdorff measure* $\mathcal{H}^s(E)$ of E is defined by

$$\mathcal{H}^s(E) := \sup_{\delta > 0} \mathcal{H}^s_\delta(E).$$

The *Hausdorff dimension* of E is defined as

$$\dim_H E := \inf\{s \geq 0 | \mathcal{H}^s(E) = 0\} = \sup\{s \geq 0 | \mathcal{H}^s(E) = \infty\}.$$

Recall the definition of packing measure, introduced by Tricot [24], Taylor & Tricot [23], which requires two limiting procedures. For $E \subset \mathbb{R}^d$ and $\delta > 0$, a *δ-packing* of E is a countable family of disjoint open balls of radii at most δ and with centers in E. For $s \geq 0$, the *s-dimensional packing premeasure* of E is defined as

$$P^s(E) := \inf_{\delta > 0}\{P^s_\delta(E)\},$$

where $P^s_\delta(E) := \sup\{\sum_i |B_i|^s\}$ with the supremum taken over all δ-packing of E. The *s-dimensional packing measure* of E is defined as

$$\mathcal{P}^s(E) := \inf\{\sum_i P^s(E_i) | E \subset \bigcup_i E_i\}.$$

The *packing dimension* of E is defined as

$$\dim_P E := \inf\{s \geq 0 | \mathcal{P}^s(E) = 0\} = \sup\{s \geq 0 | \mathcal{P}^s(E) = \infty\}.$$

Both the Hausdorff measure and dimension and the packing measure and dimension play an important role in the study of fractal geometry (see [4] and [13] for further properties of the above measures and dimensions).

2.1 Linear Cantor sets with generalized finite type condition

Let $\{S_1, \cdots, S_m\}$ be an linear IFS of contractive similitudes on the line $\mathbb{R}$ defined by

$$S_j(x) = \rho_j x + b_j, j = 1, \cdots, m, \tag{2.1}$$

with contraction ratios satisfying $0 < |\rho_j| < 1$. Let K denote the attractor of the IFS (2.1), i.e., the unique compact set $K \subset \mathbb{R}$ satisfying the equation

$$K = \bigcup_{j=1}^{m} S_j(K).$$

For each positive integer k, define the following set of finite indices

$$\Sigma_k := \{1, \cdots, m\}^k = \{(i_1, \cdots, i_k) : i_j \in \{1, \cdots, m\}, 1 \leq j \leq k\},$$

called the set of *indices of length* k with symbols $\{1, \cdots, m\}$. Also, for $k = 0$, set $\Sigma_0 := \{\emptyset\}$ and call $\emptyset$ the *empty index*. Moreover, define

$$\Sigma_* := \bigcup_{k \geq 0} \Sigma_k,$$

the *set of finite indices* with symbols $\{1, \cdots, m\}$. For $\mathbf{i} = (i_1, \cdots, i_k) \in \Sigma_*$, $E \subset \mathbb{R}$, we use the standard notation

$$S_{\mathbf{i}} := S_{i_1} \circ \cdots \circ S_{i_k}, \quad \rho_{\mathbf{i}} := \rho_{i_1} \cdots \rho_{i_k}, \quad E_{\mathbf{i}} = S_{\mathbf{i}}(E),$$

with $S_\emptyset := I$, the identity map on $\mathbb{R}$, and $\rho_\emptyset := 1$. Let $\mathbf{i} = (i_1, \cdots, i_k)$ and $\mathbf{j} = (j_1, \cdots, j_{k'})$ be two indices of Σ_*. The *length* of $\mathbf{i}$ is $|\mathbf{i}| = k$. We write $\mathbf{i} \preceq \mathbf{j}$ if $\mathbf{i}$ is an initial segment of $\mathbf{j}$, and write $\mathbf{i} \not\preceq \mathbf{j}$ if $\mathbf{i}$ is not an initial segment of $\mathbf{j}$. $\mathbf{i}$ and $\mathbf{j}$ are *incomparable* if neither $\mathbf{i} \preceq \mathbf{j}$ nor $\mathbf{j} \preceq \mathbf{i}$.

Let $\{\mathcal{M}_k\}_{k=0}^\infty$ be a sequence of index sets, where $\mathcal{M}_k \subset \Sigma_*$ for all $k \geq 0$ and $\mathcal{M}_0 = \Sigma_0$. We say that $\{\mathcal{M}_k\}_{k=0}^\infty$ is a *sequence of nested index sets* if it satisfies the following conditions:

(1) Both $\{\min\{|\mathbf{i}| : \mathbf{i} \in \mathcal{M}_k\}\}_{k=0}^{\infty}$ and $\{\max\{|\mathbf{i}| : \mathbf{i} \in \mathcal{M}_k\}\}_{k=0}^{\infty}$ are non-decreasing and have infinity limit;

(2) For each $k \geq 0$, all $\mathbf{i}, \mathbf{j} \in \mathcal{M}_k$ are incomparable if $\mathbf{i} \neq \mathbf{j}$;

(3) For each $\mathbf{j} \in \Sigma_*$ with $|\mathbf{j}| > \max\{|\mathbf{i}| : \mathbf{i} \in \mathcal{M}_k\}$, there exists $\mathbf{i} \in \mathcal{M}_k$ such that $\mathbf{i} \preceq \mathbf{j}$;

(4) For each $\mathbf{j} \in \Sigma_*$ with $|\mathbf{j}| < \min\{|\mathbf{i}| : \mathbf{i} \in \mathcal{M}_k\}$, there exists $\mathbf{i} \in \mathcal{M}_k$ such that $\mathbf{j} \preceq \mathbf{i}$;

(5) There exists a positive integer L such that for all $\mathbf{i} \in \mathcal{M}_k$ and $\mathbf{j} \in \mathcal{M}_{k+1}$ with $\mathbf{i} \preceq \mathbf{j}$, we have $|\mathbf{j}| - |\mathbf{i}| \leq L$, where L is independent of k.

Remark 2.1. If we let $\mathcal{M}_k = \Sigma_k$ for all $k \geq 0$, we get a canonical sequence of nested index sets. For general sequences, we allow $\mathcal{M}_k \cap \mathcal{M}_{k+1} \neq \emptyset$ and $\bigcup_{k=0}^{\infty} \mathcal{M}_k$ may be a proper subset of Σ_*.

The following is a non-trivial example. The sequence of nested index sets in this example is used in the original finite type condition in [18].

Example 2.2. For $k \geq 0$, let

$$\mathcal{M}_k = \Lambda_k := \{\mathbf{i} = (i_1, \cdots, i_n) \in \Sigma_* : \rho_{\mathbf{i}} \leq \rho_{\min}^k < \rho_{i_1, \cdots, i_{n-1}}\}.$$

It is easy to see that conditions $(1) - (5)$ hold by taking $L = [\ln \rho_{\min} / \ln \rho_{\max}] + 1$ in condition (5), where $\rho_{\min} := \min\{\rho_j : 1 \leq j \leq m\}$ and $\rho_{\max} := \max\{\rho_j : 1 \leq j \leq m\}$.

Note that if $\mathcal{O} \subset \mathbb{R}$ is a non-empty bounded open set which is *invariant* under $\{S_j\}_{j=1}^m$, i.e., $\bigcup_{j=1}^m \mathcal{O}_j \subset \mathcal{O}$, $\{\bigcup_{\mathbf{i} \in \mathcal{M}_k} \mathcal{O}_{\mathbf{i}}\}_{k=0}^{\infty}$ is a sequence of nested subsets of $\mathbb{R}$.

Fix a sequence of nested index sets $\{\mathcal{M}_k\}_{k=0}^{\infty}$. For each integer $k \geq 0$, let $\mathcal{V}_k$ be the set of *vertices* defined as

$$\mathcal{V}_k := \{(S_{\mathbf{i}}, k) : \mathbf{i} \in \mathcal{M}_k\}.$$

We call $(I, 0)$ the *root vertex* and let $\mathcal{V} := \bigcup_{k \geq 0} \mathcal{V}_k$. Note that if $S_{\mathbf{i}} = S_{\mathbf{j}}$ for some $\mathbf{i} \neq \mathbf{j} \in \mathcal{M}_k$, they determine the same vertex. We use the following notation for simplicity. For $\mathbf{v} = (S_{\mathbf{i}}, k) \in \mathcal{V}_k$, we introduce the convenient notation $S_{\mathbf{v}} := S_{\mathbf{i}}$ and $\rho_{\mathbf{v}} := \rho_{\mathbf{i}}$. The notation $S_{\mathbf{v}}$ allows us to refer to a vertex in $\mathcal{V}_k$ without explicitly specifying the index $\mathbf{i}$.

For simplicity, we assume that there is a following form of invariant set which will be used in the description of the generalized finite type condition: an open interval $\mathcal{O}$ which is invariant under $\{S_j\}_{j=1}^m$. (There are examples where the generalized finite type condition holds, but not with an open interval.) Without loss of generality we take $\mathcal{O} = (0, 1)$.

For any $k \geq 0$, let $F_k = \bigcup_{\mathbf{v} \in \mathcal{V}_k} \overline{\mathcal{O}}_{\mathbf{v}}$. Notice that $\overline{\mathcal{O}}_{\mathbf{v}} \subset \overline{\mathcal{O}}$, then we have

$$K = \bigcap_{k=0}^{\infty} F_k.$$

For each $\mathbf{v} \in \mathcal{V}_k$, $\overline{\mathcal{O}}_{\mathbf{v}}$ is a sub-interval contained in $[0, 1]$ with endpoints $S_{\mathbf{v}}(0)$ and $S_{\mathbf{v}}(1)$. We call $\overline{\mathcal{O}}_{\mathbf{v}}$ a *k-th generation interval* of K.

We call two intervals are *separate* if they have at most one common point. Otherwise, we call they are *overlapping*. From the definition of F_k, we see that $\bigcup_{\mathbf{v}\in\mathcal{V}_k}\mathcal{O}_{\mathbf{v}}$ consists of some separate open intervals, and each open interval is an union of the interiors of one or several k-th generation intervals. We call the closure of each such open interval a *k-th generation island*, and use $\mathcal{F}_k^0$ to denote the set of all k-th generation islands, and $\mathcal{F}_k$ the finite field generated from $\mathcal{F}_k^0$. We call the unique element $\overline{\mathcal{O}}=[0,1]$ in $\mathcal{F}_0^0$ the *root island*. For the open intervals between each pair of the k-th generation islands, we call *lakes*. For each k-th generation island I, we use $V(I)$ to denote the vertices set of all k-th generation intervals contained in I, i.e.,

$$V(I)=\{\mathbf{v}\in\mathcal{V}_k:\overline{\mathcal{O}}_{\mathbf{v}}\subset I\}.$$

It is easy to verify that $I=\bigcup_{\mathbf{v}\in V(I)}\overline{\mathcal{O}}_{\mathbf{v}}$. We call each such interval $\overline{\mathcal{O}}_{\mathbf{v}}$ a *constitutive interval* of I. Let $I\in\mathcal{F}_k^0$ and $I'\in\mathcal{F}_{k+1}^0$ for some $k\geq 0$. Then either $I'\subset I$ or they are separate. If it is in the first case, we call I a *parent* of I' and I' an *offspring*(or *descendant*) of I.

We define an equivalence relation on $\mathcal{F}^0:=\bigcup_{k\geq 0}\mathcal{F}_k^0$ to identify islands that are isomorphic in the sense that they behave the same overlap type.

Definition 2.3. Two islands $I\in\mathcal{F}_k^0$ and $I'\in\mathcal{F}_{k'}^0$ are *equivalent*, denoted by $I\sim I'$, if, there is a linear function τ mapping I onto I', such that the following conditions are satisfied:

(1) $\{S_{\mathbf{v}'}:\mathbf{v}'\in V(I')\}=\{\tau\circ S_{\mathbf{v}}:\mathbf{v}\in V(I)\}$;

(2) For any positive integer $l\geq 1$, there is an island $J\in\mathcal{F}_{k+l}^0$ contained in I if and only if there is also an island $J'\in\mathcal{F}_{k'+l}^0$ contained in I' where $J'=\tau(J)$.

It is easy to see that $\sim$ is an equivalence relation. We denote the equivalence class containing I by $[I]$ and call it the *overlap type* of I. Condition (2) says that any two islands with the same overlap type have equivalent offsprings.

Definition 2.4. We say that an linear IFS of contractive similitudes on $\mathbb{R}$ is of *generalized finite type*, or that it satisfies the *generalized finite type condition*, with respect to the invariant set $\mathcal{O}=(0,1)$, if $\mathcal{O}$ is an invariant set under the IFS and there is a sequence of nested index sets $\{\mathcal{M}_k\}_{k=0}^{\infty}$, such that $\mathcal{F}^0/\sim=\{[I]:I\in\mathcal{F}^0\}$ is a finite set.

It is easy to see that the IFS is of generalized finite type if and only if there exists some $k_0\geq 0$ such that none of the islands in $\mathcal{F}_{k_0+1}^0$ is of a new overlap type. Let $\mathcal{T}_1,\cdots,\mathcal{T}_q$ denote all the distinct overlap types, with $\mathcal{T}_1=[\overline{\mathcal{O}}]$. Now we show some examples.

Example 2.5. If $\{S_j\}_{j=1}^m$ satisfies the open set condition (where the open set is the open interval (0,1)), it is of the generalized finite type with respect to the invariant set $(0,1)$.

Proof. For each $k \geq 0$, let $\mathcal{M}_k = \Sigma_k$. It is easy to verify that in this example, the concept of k-th generation island coincides with the concept of k-th generation interval. For each k-th generation island I, $V(I)$ consist of exact only one vertex. We denote it by $\mathbf{v}(I)$. For every two islands $I \in \mathcal{F}_k^0$ and $I' \in \mathcal{F}_{k'}^0$, let $\tau = S_{\mathbf{v}(I')} S_{\mathbf{v}(I)}^{-1}$. Then τ, mapping I onto I', satisfies the conditions in Definition 2.3. Hence I and I' are equivalent. From the arbitrariness of I and I', $\mathcal{F}^0/\sim = \{\mathcal{T}_1\}$, where $\mathcal{T}_1$ is the overlap type of the island $[0,1]$.□

The following example from [9, 10] is an IFS of contractive similitudes whose contraction ratios are not exponentially commensurable. As proved in [9], it satisfies the generalized finite type condition, but not satisfies the finite type condition defined in [18].

Example 2.6. Let $\{S_j\}_{j=1}^3$ be an IFS on $\mathbb{R}$ as follows.

$$S_1(x) = \rho x, \quad S_2(x) = rx + \rho(1-r), \quad S_3(x) = rx + (1-r),$$

where $0 < \rho < 1$, $0 < r < 1$, and $\rho + 2r - \rho r \leq 1$. Then $\{S_j\}_{j=1}^3$ is of generalized finite type with respect to the invariant set $(0,1)$.



(a) (b)

Figure 1. (a) The first five levels of islands in Example 2.6 with parameters $\rho = 1/3$ and $r = 1/4$. Note that vertical bars have been added at island boundaries. (b) How the distinct overlap types are generated in Example 2.6. Iterates of the point 0 under the similitudes are represented by dots.

Proof. For each $k \geq 0$, let $\mathcal{M}_k = \Sigma_k$. Upon iterating the IFS once, the root island $I_0 = [0,1]$ generates two first generation islands:

$$I_1 = S_1([0,1]) \cup S_2([0,1]), \quad I_2 = S_3([0,1]).$$

Obviously, since $\sharp V(I_2)=1$, it follows that $I_2 \sim I_0$ with $\tau = S_3^{-1}$. It is easy to verify that $[I_0]$ and $[I_1]$ denoted respectively by $\mathcal{T}_1$ and $\mathcal{T}_2$ are distinct overlap types. Moreover, upon one more iteration, noting that $S_{13}=S_{21}$, I_1 generates three 2-th generation islands:

$$I_3 = S_{11}([0,1]) \cup S_{12}([0,1]), \quad I_4 = S_{13}([0,1]) \cup S_{22}([0,1]), \quad I_5 = S_{23}([0,1]).$$

It is straightforward to verify that

$$[I_3]=[I_4]=\mathcal{T}_2, \quad [I_5]=\mathcal{T}_1.$$

Since no new overlap types are generated, we have $\mathcal{F}^0/\sim = \{\mathcal{T}_1, \mathcal{T}_2\}$ which yields the result. See Figure 1. $\square$

Remark 2.7. In Example 2.6, there exist touching islands if and only if the contractive ratios ρ and r satisfy the equality $\rho + 2r - \rho r = 1$. However, in this case, the invariant set K is trivial, i.e., $K=[0,1]$.

The following example from [18] is an IFS of contractive similitudes whose nested index sets sequence $\{\mathcal{M}_k\}_{k=0}^{\infty}$ is taken as showed in Example 2.2, i.e., for each $k \geq 0$, $\mathcal{M}_k = \Lambda_k$.

Example 2.8. Let $\{S_j\}_{j=1}^3$ be an IFS on $\mathbb{R}$ as follows.

$$S_1(x) = \frac{1}{3}x, \quad S_2(x) = \frac{1}{9}x + \frac{8}{27}, \quad S_3(x) = \frac{1}{3}x + \frac{2}{3}.$$

Then $\{S_j\}_{j=1}^3$ is of generalized finite type with respect to the invariant set $(0,1)$.

Proof. For each $k \geq 0$, let $\mathcal{M}_k = \Lambda_k$. First, the root island $I_0=[0,1]$ generates four first generation islands in $\mathcal{F}_1^0$:

$$I_1 = S_{11}([0,1]) \cup S_{12}([0,1]), \quad I_2 = S_{13}([0,1]) \cup S_2([0,1]),$$

$$I_3 = S_{31}([0,1]) \cup S_{32}([0,1]), \quad I_4 = S_{33}([0,1]).$$

It follows from $\sharp V(I_4)=1$, that $I_4 \sim I_0$ via the map $\tau_1 = S_{33}^{-1}$. Also $I_3 \sim I_1$ with $\tau_2 = S_1 \circ S_3^{-1}$. It is easy to verify that $[I_0]$, $[I_1]$ and $[I_2]$ denoted respectively by $\mathcal{T}_1$, $\mathcal{T}_2$ and $\mathcal{T}_3$ are distinct overlap types. Moreover, noting that $S_{1133}=S_{121}$, I_1 generates five 2-th generation islands in $\mathcal{F}_2^0$:

$$I_5 = S_{1111}([0,1]) \cup S_{1112}([0,1]), \quad I_6 = S_{1113}([0,1]) \cup S_{112}([0,1]),$$

$$I_7 = S_{1131}([0,1]) \cup S_{1132}([0,1]), I_8 = S_{1133}([0,1]) \cup S_{122}([0,1]), I_9 = S_{123}([0,1]),$$

which are of overlap types $\mathcal{T}_2, \mathcal{T}_3, \mathcal{T}_2, \mathcal{T}_2, \mathcal{T}_1$. Similarly, noting that $S_{1331}=S_{211}$, $S_{1332}=S_{212}$ and $S_{1333}=S_{213}$, I_2 generates six 2-th generation islands in $\mathcal{F}_2^0$:

$$I_{10} = S_{1311}([0,1]) \cup S_{1312}([0,1]), I_{11} = S_{1313}([0,1]) \cup S_{132}([0,1]), I_{12} = S_{1331}([0,1]) \cup S_{1332}([0,1]),$$



(a) (b)

Figure 2. (a) The first five levels of islands in Example 2.8. Note that vertical bars have been added at island boundaries. (b) How the distinct overlap types are generated in Example 2.8. Iterates of the point 0 *under the similitudes are represented by dots.*

$$I_{13} = S_{1333}([0,1]) \cup S_{22}([0,1]), I_{14} = S_{231}([0,1]) \cup S_{232}([0,1]), I_{15} = S_{233}([0,1]),$$

which are of overlap types $\mathcal{T}_2, \mathcal{T}_3, \mathcal{T}_2, \mathcal{T}_3, \mathcal{T}_2, \mathcal{T}_1$. Hence $\mathcal{T}_1, \mathcal{T}_2$ and $\mathcal{T}_3$ are all the overlap types since no new overlap types are generated. See Figure 2.□

The following is a non-trivial example which allows touching islands.

Example 2.9. Let $\{S_j\}_{j=1}^4$ be an IFS on $\mathbb{R}$ as follows.

$$S_1(x) = \frac{1}{4}x, \quad S_2(x) = \frac{1}{4}x + \frac{1}{4}, \quad S_3(x) = \frac{1}{4}x + \frac{3}{8}, \quad S_4(x) = \frac{1}{4}x + \frac{3}{4}.$$

Then $\{S_j\}_{j=1}^4$ is of generalized finite type with respect to the invariant set $(0,1)$.

Proof. For each $k \geq 0$, let $\mathcal{M}_k = \Sigma_k$. First, the root island $I_0 = [0,1]$ generates three first generation islands in $\mathcal{F}_1^0$:

$$I_1 = S_1([0,1]), \quad I_2 = S_2([0,1]) \cup S_3([0,1]), \quad I_3 = S_4([0,1]),$$

where I_1 and I_2 are two touching islands. It follows from $\sharp V(I_1) = 1$ and $\sharp V(I_3) = 1$, that $I_1 \sim I_0$ and $I_3 \sim I_0$ via the map $\tau_1 = S_1^{-1}$ and $\tau_2 = S_3^{-1}$ respectively. Obviously, $[I_0]$ and $[I_2]$ denoted respectively by $\mathcal{T}_1$, $\mathcal{T}_2$ are distinct overlap types. Moreover, noting that $S_{24} = S_{32}$, I_2 generates four 2-th generation islands in $\mathcal{F}_2^0$:

$$I_4 = S_{21}([0,1]), \quad I_5 = S_{22}([0,1]) \cup S_{23}([0,1]) \cup S_{31}([0,1]),$$

$$I_6 = S_{24}([0,1]) \cup S_{33}([0,1]), \quad I_7 = S_{34}([0,1]).$$



Figure 3. (a) The first five levels of islands in Example 2.9. Note that vertical bars have been added at island boundaries. (b) How the distinct overlap types are generated in Example 2.9. Iterates of the point 0 *under the similitudes are represented by dots.*

It is straightforward to verify that $[I_4] = [I_7] = \mathcal{T}_1$, $[I_6] = \mathcal{T}_2$ and I_5 is of a new overlap type denoted by $\mathcal{T}_3$. Similarly, noting that $S_{224} = S_{232}$ and $S_{234} = S_{312}$ one may verify that I_5 generates five 3-th generation islands in $\mathcal{F}_3^0$:

$$I_8 = S_{221}([0,1]), \quad I_9 = S_{222}([0,1]) \cup S_{223}([0,1]) \cup S_{231}([0,1]),$$

$$I_{10} = S_{224}([0,1]) \cup S_{233}([0,1]) \cup S_{311}([0,1]), I_{11} = S_{234}([0,1]) \cup S_{313}([0,1]), I_{12} = S_{314}([0,1])$$

which are of overlap types $\mathcal{T}_1, \mathcal{T}_3, \mathcal{T}_3, \mathcal{T}_2, \mathcal{T}_1$. Hence $\mathcal{T}_1, \mathcal{T}_2$ and $\mathcal{T}_3$ are all the overlap types since no new overlap types are generated. See Figure 3. □

2.2 Dimension results of linear Cantor sets

In the following we assume that the linear IFS $\{S_j\}_{j=1}^m$ is of generalized finite type with respect to the invariant set $\mathcal{O} = (0,1)$ for a fixed sequence of nested index sets $\{\mathcal{M}_k\}_{k=0}^\infty$. Let $\mathcal{T}_1, \cdots, \mathcal{T}_q$ denote all the distinct overlap types, with $\mathcal{T}_1 = [\overline{\mathcal{O}}]$. For each $\alpha \geq 0$ we define a *weighted incidence matrix* $A_\alpha = A_\alpha(i,j)_{i,j=1}^q$ as follows. Fix $i (1 \leq i \leq q)$ and an island $I \in \mathcal{F}^0$ with $[I] = \mathcal{T}_i$. Suppose that I is a k-th generation island, let $I_1, \cdots, I_l$ be the offsprings of I in $\mathcal{F}_{k+1}^0$. Then we define

$$A_\alpha(i,j) := \sum \{ (\frac{|I_s|}{|I|})^\alpha : [I_s] = \mathcal{T}_j, 1 \leq s \leq l \}.$$

It is easy to see that the definition of $A_\alpha(i,j)$ is independent of the choice of I. By a similar discussion as in [9], the common value of the Hausdorff and packing dimensions of K is given by the unique α such that the spectral radius of A_α is equal to 1.

Theorem 2.10[9]. *Let λ_α be the spectral radius of the associated weighted incidence matrix A_α. Then*

$$\dim_H K = \dim_P K = \alpha,$$

where α is the unique number such that $\lambda_\alpha = 1$. Moreover, $0 < \mathcal{H}^\alpha(K) < \infty$.

For the convenience of the readers, we would like to give a direct and elementary proof of this result. The proof makes use of the ideas in [9] and [14].

Proof of Theorem 2.10. We need to define a natural probability measure μ on K. Let $(a_1, \cdots, a_q)^T$ be a 1-eigenvector of A_α, normalized so that $a_1 = 1$. (This is possible because all overlap types are descendants of $\mathcal{T}_1$.) Here α is the unique number such that $\lambda_\alpha = 1$. For each island I, where $I \in \mathcal{F}_k^0$ and $[I] = \mathcal{T}_i$ for some $k \ge 0$ and $1 \le i \le q$, we let

$$\mu(I) = |I|^\alpha a_i.$$

Obviously, $\mu([0,1]) = a_1 = 1$.

To show that μ is indeed a probability measure on K, we notice that two islands $I \in \mathcal{F}_k^0$ and $I' \in \mathcal{F}_{k'}^0$ with $k \le k'$, are overlapping if and only if either $I = I'$ in the case $k = k'$ or I' is a descendant of I in the case $k < k'$. In both cases, $I' \subset I$. Now let $I \in \mathcal{F}_k^0$ and let $\mathcal{D}$ denote the set of all offsprings of I in $\mathcal{F}_{k+1}^0$. Then

$$\begin{aligned}
\sum_{I' \in \mathcal{D}} \mu(I') &= \sum_{j=1}^{q} \sum \{\mu(I') : I' \in \mathcal{D}, [I'] = \mathcal{T}_j\} \\
&= \sum_{j=1}^{q} \sum \{|I'|^\alpha a_j : I' \in \mathcal{D}, [I'] = \mathcal{T}_j\} \\
&= |I|^\alpha \sum_{j=1}^{q} \sum \{(\frac{|I'|}{|I|})^\alpha a_j : I' \in \mathcal{D}, [I'] = \mathcal{T}_j\} \\
&= |I|^\alpha \sum_{j=1}^{q} A_\alpha(i,j) a_j \\
&= |I|^\alpha a_i = \mu(I).
\end{aligned}$$

It follows now from $\mu([0,1]) = 1$ that μ is indeed a probability measure on K.

Lower bound. Let E be a bounded Borel subset of $\mathbb{R}$ and let $\mathcal{N}(E)$ be defined as

$$\mathcal{N}(E) := \{I \in \mathcal{F}^0 : |I| \le |E| < |\mathcal{P}(I)| \text{ and } I \cap E \ne \emptyset\},$$

where $\mathcal{P}(I)$ denotes the parent of the island I. It is easy to verify that for any bounded Borel set $E \subset \mathbb{R}$, $\sharp\mathcal{N}(E) \le C_0 := \max\{|\mathcal{P}(I)|/|I| : I \in \mathcal{F}^0\} + 2$. Note that

$$\mu(E) \le \sum_{I_j \in \mathcal{N}(E)} \mu(I_j).$$

If we assume that $[I_j] = \mathcal{T}_{i_j}$,

$$\mu(E) \le \sum_{I_j \in \mathcal{N}(E)} |I_j|^\alpha a_{i_j} \le |E|^\alpha \sum_{I_j \in \mathcal{N}(E)} a_{i_j} \le C_0 \max_{1\le i\le q} a_i |E|^\alpha.$$

Thus $\mathcal{H}^\alpha(K) > 0$ and $\dim_H K \ge \alpha$(see [4]), which is the required lower bound.

Upper bound. To obtain the upper bound $\dim_H K \le \alpha$, we first assume that A_α is irreducible and thus all the a_i's are positive. For each $k \ge 0$, $K \subset \bigcup_{I\in\mathcal{F}_k^0} I$ and

$$\begin{aligned}
\sum_{I\in\mathcal{F}_k^0} |I|^\alpha &= \sum_{i=1}^{q} \sum \{\frac{1}{a_i}|I|^\alpha a_i : I \in \mathcal{F}_k^0, [I] = \mathcal{T}_i\} \\
&\le \max_{1\le i\le q}\{\frac{1}{a_i}\} \sum_{I\in\mathcal{F}_k^0} \mu(I) \\
&= \max_{1\le i\le q}\{\frac{1}{a_i}\} < \infty.
\end{aligned}$$

Since for each $k \ge 0$, $\mathcal{F}_k^0$ is a covering of K, and $\lim_{k\to\infty} \max\{|I| : I \in \mathcal{F}_k^0\} = 0$, the definition of the Hausdorff measure implies that $\mathcal{H}^\alpha(K) < \infty$, and thus $\dim_H K \le \alpha$.

Now assume A_α is not irreducible. After an appropriate permutation we can assume that A_α has the form

$$A_\alpha = \begin{bmatrix} A_1 & * & \cdots & * \\ 0 & * & \cdots & * \\ \vdots & & \cdots & * \\ 0 & \cdots & \cdots & A_r \end{bmatrix},$$

where each A_i is either an irreducible square matrix or an 1×1 zero matrix. Let $\mathcal{E} := \{A_i : A_i \text{ is non-zero}\}$, counting multiplicity. For each A_i, let $\mathcal{T}_{A_i}$ be the collection of overlap types corresponding to A_i. For each $A_i \in \mathcal{E}$ and each island I with $[I] \in \mathcal{T}_{A_i}$, define a subset $K_{A_i}(I) \subset K$ as follows.

$$K_{A_i}(I) := \bigcap_{k'=0}^{\infty} \bigcup \{I' \in \mathcal{F}_{k+k'}^0 : [I'], [\mathcal{P}(I')], \cdots, [\mathcal{P}^{k'}(I')] \in \mathcal{T}_{A_i}, \text{ and } \mathcal{P}^{k'}(I') = I\},$$

where k is the generation number of I, i.e., $I \in \mathcal{F}_k^0$. Obviously, the proof of the irreducible case above yields $\dim_H K_{A_i}(I) \le \alpha$.

For each $A_i \in \mathcal{E}$, define

$$\mathcal{F}^0_{A_i} := \{I \in \mathcal{F}^0 : [I] \in \mathcal{T}_{A_i}\}.$$

Then it is easy to verify that

$$K = \bigcup_{A_i \in \mathcal{E}} \bigcup_{I \in \mathcal{F}^0_{A_i}} K_{A_i}(I).$$

Hence, it follows from the countable stability of the Hausdorff dimension (see [4]) that $\dim_H K \leq \alpha$, which is the required upper bound.

We have proved that $\mathcal{H}^\alpha(K) > 0$ and $\dim_H K = \alpha$. This imply that $\mathcal{H}^\alpha(K) < \infty$ since K is a self-similar set (see [4]). The proof is completed. □

Remark 2.11. From Example 2.5, the classical dimension result for K satisfying the open set condition is covered by the result of Theorem 2.10, which is also showed in [9].

Remark 2.12. The weighted incidence matrix $A_\alpha = A_\alpha(i,j)_{i,j=1}^q$ can be replaced by an analogous matrix $\widetilde{A}_\alpha = \widetilde{A}_\alpha(i,j)_{i,j=1}^q$ for each $\alpha \geq 0$. For each $1 \leq i \leq q$, fix an *initial island* I_i^0 of $\mathcal{T}_i$ type. Every such island I_i^0 can be chosen arbitrarily. For example, one may choose I_i^0 as the island of $\mathcal{T}_i$ type that has the smallest generation number. (If such island is not unique, take the one lying on the left end.) Then we define $\widetilde{A}_\alpha = \widetilde{A}_\alpha(i,j)_{i,j=1}^q$ as follows. For $1 \leq i \leq q$, take an island $I \in \mathcal{F}^0$ such that $[I] = \mathcal{T}_i$. Suppose that I is a k-th generation island. Let $I_1, \cdots, I_l$ be the offsprings of I in $\mathcal{F}^0_{k+1}$. Then we define

$$\widetilde{A}_\alpha(i,j) := \sum \{ (\frac{|I_s|/|I_j^0|}{|I|/|I_i^0|})^\alpha : [I_s] = \mathcal{T}_j, 1 \leq s \leq l\}.$$

It is easy to find that $\widetilde{A}_\alpha = P_\alpha A_\alpha P_\alpha^{-1}$ where P_α is a diagonal matrix in which the elements of the main diagonal are $P_\alpha(i,i) = |I_i^0|^\alpha$ for $1 \leq i \leq q$. Hence the new weighted incidence matrix $\widetilde{A}_\alpha$ has the same spectral radius as that of A_α for each $\alpha \geq 0$. At first glance, it seems that there is almost no difference between the two kinds of matrices. However, the revised matrices $\widetilde{A}_\alpha$ are easier to calculate than the original ones. One can see this in the following examples.

Example 2.13. Let K be the invariant set of the IFS $\{S_j\}_{j=1}^3$ on $\mathbb{R}$ defined in Example 2.6. Then the common value of the Hausdorff and packing dimensions of K is α, where α is the unique solution of the equation

$$\rho^\alpha + 2r^\alpha - (\rho r)^\alpha = 1.$$

Moreover, $0 < \mathcal{H}^\alpha(K) < \infty$.

We remark that this result has already been obtained in [10] and reproved in [9].

Proof of Example 2.13. We adopt the same setup and notations of Example 2.6, with $\mathcal{T}_1 = [I_0]$ and $\mathcal{T}_2 = [I_1]$ where $I_0 = [0,1]$ is the root island and $I_1 = S_1([0,1]) \cup S_2([0,1])$. By using the proof there and after a detailed calculation we have

$$A_\alpha = \begin{bmatrix} r^\alpha & (\rho + r - \rho r)^\alpha \\ \frac{r^{2\alpha}}{(\rho + r - \rho r)^\alpha} & \rho^\alpha + r^\alpha \end{bmatrix}.$$

Setting the spectral radius of A_α equal to 1 yields the desired result. As showed in Remark 2.12, if we choose I_0 and I_1 as the initial islands of $\mathcal{T}_1$ and $\mathcal{T}_2$ types, we will get another weighted incidence matrix

$$\widetilde{A}_\alpha = \begin{bmatrix} r^\alpha & 1 \\ r^{2\alpha} & \rho^\alpha + r^\alpha \end{bmatrix},$$

which has the same spectral radius as that of A_α for each $\alpha \geq 0$. $\square$

Example 2.14. Let K be the invariant set of the IFS $\{S_j\}_{j=1}^3$ on $\mathbb{R}$ defined in Example 2.8. Then the common value of the Hausdorff and packing dimensions of K is α, where $\alpha \approx 0.7369$ is the logarithmic ratio of the largest root of the polynomial equation

$$x^3 - 6x^2 + 5x - 1 = 0$$

to 9. Moreover, $0 < \mathcal{H}^\alpha(K) < \infty$.

We remark that the result of this example has been obtained in [18] as a example of self-similar sets satisfying the finite type condition.

Proof of Example 2.14. We adopt the same setup and notations of Example 2.8, with $\mathcal{T}_1 = [I_0]$, $\mathcal{T}_2 = [I_1]$ and $\mathcal{T}_3 = [I_2]$ where $I_0 = [0,1]$ is the root island, $I_1 = S_{11}([0,1]) \cup S_{12}([0,1])$ and $I_2 = S_{13}([0,1]) \cup S_2([0,1])$. By using the proof there and after a detailed calculation we have

$$A_\alpha = \begin{bmatrix} \frac{1}{9^\alpha} & \frac{2 \cdot 11^\alpha}{81^\alpha} & \frac{5^\alpha}{27^\alpha} \\ \frac{1}{11^\alpha} & \frac{3}{9^\alpha} & \frac{5^\alpha}{33^\alpha} \\ \frac{1}{15^\alpha} & \frac{3 \cdot 11^\alpha}{135^\alpha} & \frac{2}{9^\alpha} \end{bmatrix},$$

or(choosing I_0, I_1 and I_2 as the initial islands of $\mathcal{T}_1$ $\mathcal{T}_2$ and $\mathcal{T}_3$ types)

$$\widetilde{A}_\alpha = \begin{bmatrix} \frac{1}{9^\alpha} & 2 & 1 \\ \frac{1}{81^\alpha} & \frac{3}{9^\alpha} & \frac{1}{9^\alpha} \\ \frac{1}{81^\alpha} & \frac{3}{9^\alpha} & \frac{2}{9^\alpha} \end{bmatrix}.$$

Setting the spectral radius of A_α or $\widetilde{A}_\alpha$ equal to 1 yields the desired result. $\square$

Example 2.15. Let K be the invariant set of the IFS $\{S_j\}_{j=1}^4$ on $\mathbb{R}$ defined in Example 2.9. Then the common value of the Hausdorff and packing dimensions of K is α, where

$$\alpha = \log_4(5 + \sqrt{5}) - \frac{1}{2} \approx 0.9276.$$

Moreover, $0 < \mathcal{H}^\alpha(K) < \infty$.

Proof. We adopt the same setup and notations of Example 2.9, with $\mathcal{T}_1 = [I_0]$, $\mathcal{T}_2 = [I_2]$ and $\mathcal{T}_3 = [I_5]$ where $I_0 = [0,1]$ is the root island, $I_2 = S_2([0,1]) \cup S_3([0,1])$ and $I_5 = S_{22}([0,1]) \cup S_{23}([0,1]) \cup S_{31}([0,1])$. By using the proof there and after a detailed calculation we have

$$A_\alpha = \begin{bmatrix} \frac{2}{4^\alpha} & \frac{3^\alpha}{8^\alpha} & 0 \\ \frac{2}{6^\alpha} & \frac{1}{4^\alpha} & \frac{1}{3^\alpha} \\ \frac{2}{8^\alpha} & \frac{3^\alpha}{16^\alpha} & \frac{2}{4^\alpha} \end{bmatrix},$$

or(choosing I_0, I_2 and I_5 as the initial islands of $\mathcal{T}_1$ $\mathcal{T}_2$ and $\mathcal{T}_3$ types)

$$\widetilde{A}_\alpha = \begin{bmatrix} \frac{2}{4^\alpha} & 1 & 0 \\ \frac{2}{16^\alpha} & \frac{1}{4^\alpha} & 1 \\ \frac{2}{64^\alpha} & \frac{1}{16^\alpha} & \frac{2}{4^\alpha} \end{bmatrix}.$$

Setting the spectral radius of A_α or $\widetilde{A}_\alpha$ equal to 1 yields the desired result. □

3 Density Theorems

Let K be a linear Cantor set satisfying the generalized finite type condition with respect to the invariant open set $\mathcal{O} = (0,1)$. In this section, we analyze the local behaviour of the Hausdorff measure and the packing measure of K. In particular, we will first prove two inequalities stated in Theorem 1.1 and Theorem 1.2, i.e.,

$$\mathcal{H}^\alpha(K \cap J) \le |J|^\alpha$$

for all intervals J contained in $[0,1]$, and

$$\mathcal{P}^\alpha(K \cap J) \ge |J|^\alpha$$

for all intervals J contained in $[0,1]$ with centers in K. Here α is the number such that the spectral radius of the weighted incidence matrix A_α is equal to 1, which is also the common value of the Hausdorff and packing dimensions of K by Theorem 2.10. We assume that the matrix A_α is irreducible throughout this section and the following ones. Actually, it is easy to verify that Example 2.5, Example 2.6, Example 2.8 and Example 2.9 all satisfy this assumption.

For a matrix A, we use $r(A)$ to denote the spectral radius of A. The following basic algebraic lemma is needed.

Lemma 3.1. *Let A be a $q \times q$ non-negative irreducible matrix with $q \ge 2$, A' be the $(q-1) \times (q-1)$ sub-matrix at the right-bottom corner of A. Then $r(A') < r(A)$.*

Proof. Since A is irreducible, there exists at least one positive number in the set $\{A(1,j) : 1 \leq j \leq q\} \cup \{A(i,1) : 1 \leq i \leq q\}$. We define a $q \times q$ matrix B in which $B(1,j) = A(1,j)/2$ for $1 \leq j \leq q$, $B(i,1) = A(i,1)/2$ for $1 \leq i \leq q$ and $B(i,j) = A(i,j)$ for $2 \leq i,j \leq q$. Obviously, B is also a non-negative irreducible matrix and $B < A$, i.e., their exists at least one coordinate (i,j) such that $B(i,j) < A(i,j)$. Hence from the well-known Perron-Frobenius Theorem, it yields that $r(B) < r(A)$. Then we get the desired result since $r(A') \leq r(B)$. □

The following lemma is useful for the proof of the two inequalities.

Lemma 3.2. S*uppose A_α is irreducible, then K can be decomposed into an union of a set K_a with a graph directed construction and an attractor K_b of a countable infinite IFS. Moreover,*

$$\dim_H K_a = \dim_P K_a < \dim_H K_b = \dim_P K_b = \dim_H K = \dim_P K,$$

and

$$\mathcal{H}^\alpha(K) = \mathcal{H}^\alpha(K_b), \quad \mathcal{P}^\alpha(K) = \mathcal{P}^\alpha(K_b).$$

Proof. For each $k \geq 1$, let $\mathcal{F}_k'^0$ be a subset contained in $\mathcal{F}_k^0$ as

$$\mathcal{F}_k'^0 = \{I \in \mathcal{F}_k^0 : [I] \neq \mathcal{T}_1\}.$$

Let $K_a = \bigcap_{k\geq 1} \bigcup \{I : I \in \mathcal{F}_k'^0\}$. Then it is easy to find that K_a is a set with a graph directed construction. In fact, there are $q-1$ vertex sets in the construction of K_a whose weighted incidence matrix is the $(q-1) \times (q-1)$ sub-matrix A'_α at the right-bottom corner of A_α for each $\alpha \geq 0$. Moreover, by a similar proof of Theorem 2.10, the Hausdorff dimension of K_a is the unique α such that the spectral radius of A'_α is equal to 1 (or see a direct result in [14]). It yields from Lemma 3.1, $\dim_H K_a < \dim_H K$.

For each $I \in \mathcal{F}^0 \setminus \mathcal{F}_0^0$ with overlap type $\mathcal{T}_1$, suppose the generation number of I is k, i.e., $I \in \mathcal{F}_k^0$ with $k \geq 1$. If none of the elements in $\{\mathcal{P}(I), \cdots, \mathcal{P}^{k-1}(I)\}$ is of $\mathcal{T}_1$ type, we call I a $\mathcal{T}_1$ *type utmost island*.

For each $\mathcal{T}_1$ type utmost island I, the vertex set $V(I)$ of I consists of exactly one vertex, i.e., $\sharp V(I) = 1$. Denote the contractive similitude of the unique element in $V(I)$ as S_I. Then there exists a countable infinite IFS (see [15] for further properties of infinite IFS) of contractive similitudes

$$\mathcal{S} := \{S_I : I \in \mathcal{F}^0 \setminus \mathcal{F}_0^0, I \text{ is a } \mathcal{T}_1 \text{ type utmost island}\}.$$

Denote by K_b the attractor of $\mathcal{S}$. From the construction of K_a and K_b, one can easily observe that $K = K_a \cup K_b$. Hence, it follows from the stability property of the Hausdorff dimension (see [4]), $\dim_H K_b = \dim_H K$. The remaining is obvious. □

Remark 3.3. It is worth while to point out that in the proof of Lemma 3.2 we could replace $\mathcal{T}_1$ by any other overlap type. With appropriate modifications, we can also prove Lemma 3.2 in a similar way. We will not go into the details here.

In order to prove Theorem 1.1 and Theorem 1.2, we need a detailed analysis of the attractor K_b of a countable infinite IFS $\mathcal{S}$ described in Lemma 3.2. Now we introduce some notations for convenience. Denote the list of countable contractive similitudes in $\mathcal{S}$ as

$$\mathcal{S} = \{S'_1, \cdots, S'_j, \cdots\}$$

and r'_j the contractive ratio of S'_j. Then for each $k \geq 1$ we will write $S'_{\mathbf{i}} = S'_{i_1} \circ \cdots \circ S'_{i_k}$ and $r'_{\mathbf{i}} = r'_{i_1} \cdots r'_{i_k}$ for all indices $\mathbf{i} = i_1 \cdots i_k$ with entries $i_j \in \mathbb{N}$. Also, for every such indice $\mathbf{i} = i_1 \cdots i_k$ we will write $|\mathbf{i}| = k$ for the length of $\mathbf{i}$. From the construction of K_b, we know the intervals $S'_i([0,1])$ and $S'_j([0,1])$ are separate for all $i \neq j \in \mathbb{N}$. Hence

$$\mathcal{H}^\alpha(K_b) = \sum_{i\in\mathbb{N}} \mathcal{H}^\alpha(S'_i K_b) = \sum_{i\in\mathbb{N}} r'^\alpha_i \mathcal{H}^\alpha(K_b).$$

Combing the above formula with the result in Lemma 3.2 and the fact that $0 < \mathcal{H}^\alpha(K) < \infty$ from Theorem 2.10, we have

$$\sum_{i\in\mathbb{N}} r'^\alpha_i = 1.$$

Moreover,

$$\sum_{|\mathbf{i}|=k} r'^\alpha_{\mathbf{i}} = (\sum_{i\in\mathbb{N}} r'^\alpha_i)^k = 1$$

for any integer $k \geq 1$.

A useful measure λ on K, which is called α-dimensional normalized Hausdorff measure, defined on K_b by $\lambda(S'_{\mathbf{i}}(K_b)) = r'^\alpha_{\mathbf{i}}$, then extended to Borel subsets of K, will be used. This is a probability measure which scales on $S'_{\mathbf{i}}(K_b)$, hence $\mathcal{H}^\alpha|_{K_b} = \mathcal{H}^\alpha|_K = \mathcal{H}^\alpha(K)\lambda$ by Lemma 3.2 and the scaling property of the Hausdorff measure on K_b. After a similar discussion, one can also get $\mathcal{P}^\alpha|_{K_b} = \mathcal{P}^\alpha|_K = \mathcal{P}^\alpha(K)\lambda$.

Remark 3.4. We should point out that the α-dimensional normalized Hausdorff measure λ is equal to the natural probability measure μ defined in the proof of Theorem 2.10. This could be easily verified by showing λ and μ are equal on cylinder sets in K_b, i.e.,

$$\mu(S'_{\mathbf{i}}(K_b)) = |S'_{\mathbf{i}}([0,1])|^\alpha a_1 = r'^\alpha_{\mathbf{i}} = \lambda(S'_{\mathbf{i}}(K_b)),$$

where $a_1 = 1$ is the first element in the normalized 1-eigenvector of A_α as introduced in the proof of Theorem 2.10. Hence for each island $I \in \mathcal{F}^0$, $\lambda(I)$ can be calculated by using the parameters of the IFS of K.

For any interval $J \subset [0,1]$ we define the *density* of J with respect to λ as

$$d(J) = \frac{\lambda(J)}{|J|^\alpha}.$$

Theorem 1.1 and Theorem 1.2 are used to obtain density results for Hausdorff and packing measures of K, which can be further used to compute the exact value of the α-dimensional Hausdorff measure $\mathcal{H}^\alpha(K)$ of K and the exact value of the α-dimensional packing measure $\mathcal{P}^\alpha(K)$ of K. These are the following formulae

$$\mathcal{H}^\alpha(K)^{-1} = \sup\{d(J) : J \text{ is an interval with } J \subset [0,1]\},$$

and

$$\mathcal{P}^\alpha(K)^{-1} = \inf\{d(J) : J \text{ is an interval centered in } K \text{ with } J \subset [0,1]\},$$

which are stated in Corollary 1.3 and Corollary 1.4.

The above two formulae are useful for computing $\mathcal{H}^\alpha(K)$ and $\mathcal{P}^\alpha(K)$ for the following reason. Remark 3.4 ensures us to make a detailed analysis of λ, and thus a possible detailed analysis of the supremum and infimum in the above two explicit formulae.

3.1 Proof of Theorem 1.1 and Corollary 1.3

We analyze the local behaviour of the α-dimensional Hausdorff measure $\mathcal{H}^\alpha(K)$ of K in this subsection. The principal result in this subsection, i.e., Theorem 1.1 states that

$$\mathcal{H}^\alpha(K \cap J) \leq |J|^\alpha$$

for all intervals $J \subset [0,1]$. This formula is analogous to that of a general self-similar set in $\mathbb{R}^d$ which satisfies the open set condition (see [19]). Using Lemma 3.2, it is now possible to adapt the techniques for proving the analogous formula in [19] to establish Theorem 1.1.

We turn to show two more lemmas concerning the α-dimensional Hausdorff measure $\mathcal{H}^\alpha(K_b)$ of the attractor K_b as follows. Theorem 1.1 will be a direct corollary of them.

Lemma 3.5. *Let K_b be the attractor described in Lemma 3.2, $J \subset [0,1]$ be an interval and k be a positive integer, then*

$$\mathcal{H}^\alpha(K_b \cap \bigcup_{|\boldsymbol{i}|=k} S'_{\boldsymbol{i}} J) \geq \mathcal{H}^\alpha(K_b \cap J).$$

Proof. Since $S'_{\mathbf{j}} K_b \subset \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} K_b = K_b$ for all $\mathbf{j}$ with $|\mathbf{j}| = k$, we have

$$K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J \supset \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}(K_b \cap J).$$

It follows that

$$\begin{aligned}\mathcal{H}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) &\geq \mathcal{H}^\alpha(\bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}(K_b \cap J)) \\ &= \sum_{|\mathbf{i}|=k} \mathcal{H}^\alpha(S'_{\mathbf{i}}(K_b \cap J)) \\ &= \sum_{|\mathbf{i}|=k} r'^{\alpha}_{\mathbf{i}} \mathcal{H}^\alpha(K_b \cap J) \\ &= \mathcal{H}^\alpha(K_b \cap J). \quad \Box\end{aligned}$$

The following lemma is a revised form of Theorem 1.1 with K replaced by K_b.

Lemma 3.6. *The attractor K_b described in Lemma 3.2 satisfies*

$$\mathcal{H}^\alpha(K_b \cap J) \leq |J|^\alpha \tag{3.1}$$

for all intervals $J \subset [0,1]$.

Proof. In order to reach a contradiction, we assume that (3.1) is not satisfied, i.e., there exists a non-empty interval $J \subset [0,1]$, such that

$$\mathcal{H}^\alpha(K_b \cap J) > |J|^\alpha.$$

It follows from this we can find $0 < \kappa < 1$ with

$$(1-\kappa)\mathcal{H}^\alpha(K_b \cap J) > |J|^\alpha. \tag{3.2}$$

Next, fix $\delta > 0$ and choose a positive integer k such that $|S'_{\mathbf{i}} J| \leq \delta$ for all $\mathbf{i}$ with $|\mathbf{i}| = k$. Let

$$\eta = \frac{1}{2}\kappa \mathcal{H}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J).$$

It follows from Lemma 3.5 and (3.2),

$$\mathcal{H}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) \geq \mathcal{H}^\alpha(K_b \cap J) \geq \frac{|J|^\alpha}{1-\kappa} > 0, \tag{3.3}$$

which yields $\eta > 0$.

Since $\eta > 0$, we can choose a covering $\{J_i\}_i$ of $K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J$ with $|J_i| \leq \delta$ such that

$$\begin{aligned}\sum_i |J_i|^\alpha &\leq \mathcal{H}^\alpha_\delta(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) + \eta \\ &\leq \mathcal{H}^\alpha(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) + \eta \end{aligned} \tag{3.4}$$

The family $\{S'_{\mathbf{i}}J\}_{|\mathbf{i}|=k} \cup \{J_i\}_i$ is clearly a δ-covering of K_b. We therefore conclude from (3.2), (3.4) and Lemma 3.5 that

$$\begin{aligned}
\mathcal{H}^{\alpha}_{\delta}(K_b) &\le \sum_{|\mathbf{i}|=k} |S'_{\mathbf{i}}J|^{\alpha} + \sum_{i} |J_i|^{\alpha} \\
&\le \sum_{|\mathbf{i}|=k} r'^{\alpha}_{\mathbf{i}} |J|^{\alpha} + \mathcal{H}^{\alpha}(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) + \eta \\
&= |J|^{\alpha} + \mathcal{H}^{\alpha}(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) + \eta \\
&\le (1-\kappa)\mathcal{H}^{\alpha}(K_b \cap J) + \mathcal{H}^{\alpha}(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) + \eta \\
&\le (1-\kappa)\mathcal{H}^{\alpha}(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) + \mathcal{H}^{\alpha}(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) + \eta \\
&\le \mathcal{H}^{\alpha}(K_b) - \kappa\mathcal{H}^{\alpha}(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) + \eta \\
&= \mathcal{H}^{\alpha}(K_b) - \eta \\
&\le \mathcal{H}^{\alpha}(K_b) - \frac{1}{2}\mathcal{H}^{\alpha}(K_b \cap J).
\end{aligned}$$

Finally, letting $\delta \to 0$ gives

$$\mathcal{H}^{\alpha}(K_b) \le \mathcal{H}^{\alpha}(K_b) - \frac{1}{2}\mathcal{H}^{\alpha}(K_b \cap J).$$

Then the fact that $0 < \mathcal{H}^{\alpha}(K_b) < \infty$ by Lemma 3.2 and $(1/2)\mathcal{H}^{\alpha}(K_b \cap J) > 0$ by (3.3) provides the desired contradiction. □

Proof of Theorem 1.1. Indeed, it follows from Lemma 3.2 and Lemma 3.6. □

Putting J equal to $[0,1]$ in Theorem 1.1 gives the upper bound for the α-dimensional Hausdorff measure of K, namely, $\mathcal{H}^{\alpha}(K) \le 1$. It is a natural generalization of the same result in the open set condition case, since in that case K can be covered by its iterated images under the IFS. Recall that it is not always true that we have equality. See Falconer [4] for some examples with $\mathcal{H}^{\alpha}(K) = 1$ satisfying the open set condition. An natural question is arisen: *is there any non-trivial example with* $\mathcal{H}^{\alpha}(K) = 1$ *which satisfies only the generalized finite type condition.* Moreover, Theorem 1.1 has an application on densities and can be applied to compute the exact value of the Hausdorff measure $\mathcal{H}^{\alpha}(K)$ of K. We will now elaborate on this.

For a given measure ν on $\mathbb{R}$ and $s > 0$, the *upper s-dimensional convex density* of ν at x is defined by

$$\mathcal{D}^{*s}(\nu, x) = \lim_{r\to 0} \sup\{\frac{\nu(J)}{|J|^s} : J \text{ is an interval and } 0 < |J| \le r, x \in J\}.$$

The *lower s-dimensional convex density* $\mathcal{D}^s_*(\nu, x)$ is defined similarly by taking the lower limit. We have the following result that if $E \subset \mathbb{R}$ and $s > 0$ with $\mathcal{H}^s(E) < \infty$,

$$\mathcal{D}^{*s}(\mathcal{H}^s|_E, x) = 1 \text{ for } \mathcal{H}^s\text{-a.e. } x \in E. \tag{3.5}$$

The reader is referred to [2] for a proof of (3.5).

We then could give the following proof of Corollary 1.3 on the basis of (3.5) and Theorem 1.1.

Proof of Corollary 1.3. From (3.5), we can pick a point $x \in K \cap (0, 1)$ such that $\mathcal{D}^{*\alpha}(\mathcal{H}^\alpha|_K, x) = 1$. By the definition of $\mathcal{D}^{*\alpha}(\mathcal{H}^\alpha|_K, x)$, there exists a positive sequence $\{\delta_n\}_n$ with $\delta_n < \min\{x, 1 - x\}$ and $\delta_n \to 0$ as $n \to \infty$ such that

$$\sup_{0<|J|\le\delta_n} \frac{\mathcal{H}^\alpha(K \cap J)}{|J|^\alpha} - \frac{1}{n} \le 1 \le \sup_{0<|J|\le\delta_n} \frac{\mathcal{H}^\alpha(K \cap J)}{|J|^\alpha} + \frac{1}{n}.$$

Hence there exists an interval J_n with $0 < |J_n| \le \delta_n$ for each n such that

$$\sup_{0<|J|\le\delta_n} \frac{\mathcal{H}^\alpha(K \cap J)}{|J|^\alpha} \le \frac{\mathcal{H}^\alpha(K \cap J_n)}{|J_n|^\alpha} + \frac{1}{n}.$$

Thus

$$\frac{\mathcal{H}^\alpha(K \cap J_n)}{|J_n|^\alpha} - \frac{1}{n} \le 1 \le \frac{\mathcal{H}^\alpha(K \cap J_n)}{|J_n|^\alpha} + \frac{2}{n},$$

which yields that

$$\frac{\mathcal{H}^\alpha(K \cap J_n)}{|J_n|^\alpha} \to 1$$

as $n \to \infty$. Moreover, by Theorem 1.1, for each interval J_n we have $\mathcal{H}^\alpha(K \cap J_n)/|J_n|^\alpha \le 1$. Hence

$$\sup\{\frac{\mathcal{H}^\alpha(K \cap J)}{|J|^\alpha} : J \subset [0, 1]\} = 1.$$

Since $\lambda = \mathcal{H}^\alpha|_K/\mathcal{H}^\alpha(K)$, (1.9) follows immediately from the above equation. □

As mentioned earlier, a special case of Theorem 1.1 and Corollary 1.3 is implicit in earlier work by Marion ([11, 12]) and Ayer & Strichartz ([1]) since the generalized finite condition with respect to the invariant open set $(0, 1)$ is a natural generalization of the open set condition with the open set $(0, 1)$. In another direction, Olsen [19] generalized Marion and Ayer & Strichartz's works to general self-similar sets in $\mathbb{R}^d$ satisfying the open set condition. With suitable modifications if necessary, we may generalize Theorem 1.1 and Corollary 1.3 in an analogous way to general self-similar sets satisfying the generalized finite type condition with irreducible weighted incidence matrix. However, in order to match the main goal of this paper and to avoid additional technical details, we will not pursue this here.

3.2 Proof of Theorem 1.2 and Corollary 1.4

In this subsection we will analyze the behaviour of the packing measure of the linear Cantor set K. Our main result in this subection, i.e., Theorem 1.2 says that

$$\mathcal{P}^{\alpha}(K \cap J) \geq |J|^{\alpha}$$

for all intervals $J \subset [0,1]$ centered in K. In a manner dual to the Hausdorff measure case, the above result is also analogous to that of a general self-similar set in $\mathbb{R}^d$ which satisfies the strong separation condition or the open set condition (see [19, 20]). Hence it is also possible to adapt the techniques for proving the analogous formulae in [19, 20] to establish Theorem 1.2 by using Lemma 3.2.

In order to prove Theorem 1.2, we shall need the following two lemmas concerning the packing measure $\mathcal{P}^{\alpha}(K_b)$ of the attractor K_b.

Lemma 3.7. *Let K_b be the attractor described in Lemma 3.2. Let $J \subset [0,1]$ be an interval centered in K_b and k a positive integer. Then*

$$\mathcal{P}^{\alpha}(K_b \cap \bigcup_{|\boldsymbol{i}|=k} S'_{\boldsymbol{i}}J) = \mathcal{P}^{\alpha}(K_b \cap J) > 0.$$

Proof. We write J° as the interior of the interval J. First, we prove that

$$K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J^{\circ} = \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}(K_b \cap J^{\circ}). \tag{3.6}$$

Fix a point $y \in K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J^{\circ}$. Since $J^{\circ} \subset \mathcal{O} = (0,1)$, there exists an index $\mathbf{u}$ with the length k such that $y \in S'_{\mathbf{u}}J^{\circ} \subset S'_{\mathbf{u}}\mathcal{O}$. We also have $y \in K_b = \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}K_b$ and we therefore find an index $\mathbf{v}$ with the length k such that $y \in S'_{\mathbf{v}}K_b \subset S'_{\mathbf{v}}\overline{\mathcal{O}}$. Thus $y \in S'_{\mathbf{u}}\mathcal{O} \cap S'_{\mathbf{v}}\overline{\mathcal{O}}$, and therefore $\mathbf{u} = \mathbf{v}$. Hence $y \in S'_{\mathbf{u}}J^{\circ} \cap S'_{\mathbf{u}}K_b = S'_{\mathbf{u}}(K_b \cap J^{\circ}) \subset \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}(K_b \cap J^{\circ})$ which yields that $K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J^{\circ} \subset \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}(K_b \cap J^{\circ})$. The other direction is obvious. Hence the formula (3.6) holds.

It follows from (3.6) that

$$\begin{aligned}
\mathcal{P}^{\alpha}(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) &= \mathcal{P}^{\alpha}(\bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}(K_b \cap J^{\circ})) \\
&= \sum_{|\mathbf{i}|=k} \mathcal{P}^{\alpha}(S'_{\mathbf{i}}(K_b \cap J^{\circ})) \\
&= \sum_{|\mathbf{i}|=k} r'^{\alpha}_{\mathbf{i}} \mathcal{P}^{\alpha}(K_b \cap J^{\circ}) \\
&= \mathcal{P}^{\alpha}(K_b \cap J).
\end{aligned}$$

Moreover, since J has its center in K_b, we deduce that $\mathcal{P}^\alpha(K_b \cap J) > 0$. This completes the proof of Lemma 3.7. $\square$

Lemma 3.8. The attractor K_b described in Lemma 3.2 satisfies

$$\mathcal{P}^\alpha(K_b \cap J) \geq |J|^\alpha \tag{3.7}$$

for all intervals $J \subset [0,1]$ centered in K_b.

Proof. In order to reach a contradiction, we assume that (3.7) is not satisfied, i.e., there exists an interval $J := [c, d] \subset [0,1]$ centered in K_b, such that

$$\mathcal{P}^\alpha(K_b \cap J) < |J|^\alpha.$$

Thus we can find a number $0 < \kappa < 1$ with

$$(1+\kappa)\mathcal{P}^\alpha(K_b \cap J) < |J|^\alpha. \tag{3.8}$$

Next, fix $\delta > 0$ and choose a positive integer k such that

$$|S'_{\mathbf{i}} J| \leq \delta$$

for all $\mathbf{i}$ with $|\mathbf{i}| = k$. Let

$$\eta = \frac{1}{2}\kappa \mathcal{P}^\alpha(K_b \cap J).$$

It follows from Lemma 3.7, $\eta > 0$.

For a positive integer n, write $G_n = K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}((c-1/n, d+1/n))$, and observe that

$$G_1 \subset G_2 \subset \cdots \subset G_n \subset \cdots$$

and

$$\bigcup_n G_n = K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J.$$

If $K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J \neq \emptyset$, there is a positive integer n_0 with $1/n_0 < \delta$ such that $G_{n_0} \neq \emptyset$, and

$$\mathcal{P}^\alpha(G_{n_0}) \geq \mathcal{P}^\alpha(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) - \frac{\eta}{2}. \tag{3.9}$$

We can choose a $1/n_0$-packing $\{J_i^\circ\}_i$ of G_{n_0} such that

$$\begin{aligned}\sum_i |J_i|^\alpha &\geq P^\alpha_{\frac{1}{n_0}}(G_{n_0}) - \frac{\eta}{2} \\ &\geq P^\alpha(G_{n_0}) - \frac{\eta}{2} \\ &\geq \mathcal{P}^\alpha(G_{n_0}) - \frac{\eta}{2}. \end{aligned} \tag{3.10}$$

Since $J^\circ \subset \mathcal{O}$, $S'_{\mathbf{i}}(J^\circ) \cap S'_{\mathbf{j}}(J^\circ) = \emptyset$ for all $\mathbf{i} \neq \mathbf{j}$ with lengths k. And for each such $\mathbf{i}$, since $(c+d)/2 \in K_b$, we have $S'_{\mathbf{i}}((c+d)/2) \in S'_{\mathbf{i}} K_b \subset K_b$. Thus the family $\{S'_{\mathbf{i}}(J^\circ)\}_{|\mathbf{i}|=k}$ is a δ-packing of $K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J$ by the fact that $|S'_{\mathbf{i}} J| \leq \delta$ for all $\mathbf{i}$ with $|\mathbf{i}| = k$.

Since $\{J_i^\circ\}_i$ is also a $1/n_0$-packing of G_{n_0}, we conclude that $\{S'_{\mathbf{i}} J^\circ\}_{|\mathbf{i}|=k} \bigcup \{J_i^\circ\}_i$ is a δ-packing of K_b. Using this we therefore conclude from (3.8), (3.9), (3.10), and Lemma 3.7 that

$$\begin{aligned}
P_\delta^\alpha(K_b) &\geq \sum_{|\mathbf{i}|=k} (r'_{\mathbf{i}}|J|)^\alpha + \sum_i (|J_i|)^\alpha \\
&\geq \sum_{|\mathbf{i}|=k} r'^{\alpha}_{\mathbf{i}} |J|^\alpha + \mathcal{P}^s(G_{n_0}) - \frac{\eta}{2} \\
&\geq |J|^\alpha + \mathcal{P}^\alpha(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) - \eta \\
&\geq (1+\kappa)\mathcal{P}^\alpha(K_b \cap J) + \mathcal{P}^\alpha(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) - \eta \\
&= (1+\kappa)\mathcal{P}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) + \mathcal{P}^\alpha(K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) - \eta \\
&= \mathcal{P}^\alpha(K_b) + \kappa \mathcal{P}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J) - \eta \\
&= \mathcal{P}^\alpha(K_b) + 2\eta - \eta \\
&= \mathcal{P}^\alpha(K_b) + \frac{1}{2}\kappa \mathcal{P}^\alpha(K_b \cap J).
\end{aligned}$$

Finally, let $\delta \to 0$, we get

$$P^\alpha(K_b) \geq \mathcal{P}^\alpha(K_b) + \frac{1}{2}\kappa \mathcal{P}^\alpha(K_b \cap J). \tag{3.11}$$

In [5] it is proved that the packing premeasure P^α coincides with the packing measure $\mathcal{P}^\alpha$ for compact subsets with finite P^α-measure. Thus they coincide for K_b, and it follows from (3.11) that

$$\mathcal{P}^\alpha(K_b) \geq \mathcal{P}^\alpha(K_b) + \frac{1}{2}\kappa \mathcal{P}^\alpha(K_b \cap J).$$

Since $\mathcal{P}^\alpha(K_b)$ is positive and finite by Lemma 3.2, and $(1/2)\kappa\mathcal{P}^\alpha(K_b \cap J) > 0$, we get the contradiction.

On the other hand, if $K_b \setminus \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J = \emptyset$, i.e., $K_b \subset \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}} J$, the aforementioned string of inequalities simplifies to

$$\begin{aligned} P_\delta^\alpha(K_b) &\geq \sum_{|\mathbf{i}|=k} (r'_{\mathbf{i}}|J|)^\alpha \\ &= \sum_{|\mathbf{i}|=k} r'^{\alpha}_{\mathbf{i}}|J|^\alpha \\ &= |J|^\alpha \\ &> (1+\kappa)\mathcal{P}^\alpha(K_b \cap J) \\ &= (1+\kappa)\mathcal{P}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J). \end{aligned}$$

Letting $\delta \to 0$ and using the fact that $\mathcal{P}^\alpha(K_b) = P^\alpha(K_b)$ gives

$$\begin{aligned} \mathcal{P}^\alpha(K_b) &= P^\alpha(K_b) \\ &\geq (1+\kappa)\mathcal{P}^\alpha(K_b \cap \bigcup_{|\mathbf{i}|=k} S'_{\mathbf{i}}J) \\ &= (1+\kappa)\mathcal{P}^\alpha(K_b) > \mathcal{P}^\alpha(K_b). \end{aligned}$$

This provides the desired contradiction.

The proof of Lemma 3.8 is completed. □

Proof of Theorem 1.2. For any $J := [c,d] \subset [0,1]$ centered in K. If J is centered in K_b, the result follows immediately from Lemma 3.2 and Lemma 3.8. If J is not centered in K_b, i.e., $(c+d)/2 \in K_a \setminus K_b$, we claim that:

Claim: $K_b \cap (c,d) \neq \emptyset$ and for each positive integer n, there exists a point $x_n \in K_b \cap (c,d)$ such that $|x_n - (c+d)/2| \leq 1/n$.

Proof. Since $(c+d)/2 \in K_a$, for each n, there is an island $I_n \subset (c,d)$ containing the point $(c+d)/2$ with length less than $1/n$ whose overlap type is not $\mathcal{T}_1$. By the irreducible property of the weighted incidence matrix A_α, we could find a smaller $\mathcal{T}_1$ type island $\widetilde{I}_n$ contained in I_n. By the construction of K_b, $K_b \cap \widetilde{I}_n \neq \emptyset$ which yields that $K_b \cap (c,d) \neq \emptyset$. Then fix a point x_n in $K_b \cap \widetilde{I}_n$. Obviously we get $|x_n - (c+d)/2| \leq |I_n| \leq 1/n$. □

By this claim we define a sequence of intervals $\{J_n\}_n$ contained in J as

$$J_n = [x_n - \min\{x_n - c, d - x_n\}, x_n + \min\{x_n - c, d - x_n\}].$$

It is not difficult to find that for each n the interval J_n is centered in K_b, and that the left endpoint of J_n tends to c and the right endpoint of J_n tends to d as $n \to \infty$. As showed in the first case, $\mathcal{P}^\alpha(K \cap J_n) \geq |J_n|^\alpha$ for all n. Letting $n \to \infty$, we immediately get $\mathcal{P}^\alpha(K \cap J) \geq |J|^\alpha$. □

In a manner dual to the Hausdorff measure case, this result also has an application on densities which can be applied to compute the exact value of the packing measure $\mathcal{P}^\alpha(K)$ of K.

Let $s \geq 0$, for a given measure ν on $\mathbb{R}$ and $x \in \mathbb{R}$, the *lower s-dimensional density* of ν at x is defined as

$$\Theta_*^s(\nu, x) = \liminf_{r \to 0} \frac{\nu([x-r, x+r])}{(2r)^s}.$$

The *upper s-dimensional density* $\Theta^{*\alpha}(\nu, x)$ is defined similarly by taking the upper limit. We have the following result. If $E \subset \mathbb{R}$ and $s > 0$ with $0 < \mathcal{P}^s(E) < \infty$,

$$\Theta_*^s(\mathcal{P}^s|_E, x) = 1 \text{ for } \mathcal{P}^\alpha\text{-a.e.} \quad x \in E. \tag{3.12}$$

See the proof in [13]. We then could prove Corollary 1.4 on the basis of (3.12) and Theorem 1.2.

Proof of Corollary 1.4. Let $\mathcal{O} = (0,1)$. Since $K \cap \mathcal{O} \neq \emptyset$, we can take a point $y \in K \cap \mathcal{O}$. Choose $\delta > 0$ such that the open interval $(y-\delta, y+\delta)$ is contained in $\mathcal{O}$. Moreover, since $y \in K$, $\mathcal{P}^\alpha(K \cap (y-\delta, y+\delta)) > 0$. Hence from (3.12), there exists a point $z \in K \cap (y-\delta, y+\delta)$ with $\Theta_*^\alpha(\mathcal{P}^\alpha|_K, z) = 1$. By the definition of $\Theta_*^\alpha(\mathcal{P}^\alpha|_K, z)$, there exists a sequence $\{r_n\}_n$ with each $r_n < \delta - d(z,y)$ and $r_n \to 0$ as $n \to \infty$, such that $\lim_{n\to\infty} \mathcal{P}^\alpha(K \cap [z-r_n, z+r_n])/(2r_n)^\alpha = 1$. Notice that all intervals $[z-r_n, z+r_n]$ are contained in $(y-\delta, y+\delta) \subset \mathcal{O}$ with center $z \in K$. However, by Theorem 1.2, for each interval $J \subset [0,1]$ centered in K, we have $\mathcal{P}^\alpha(K \cap J)/|J|^\alpha \geq 1$. Hence we get

$$\inf\{\frac{\mathcal{P}^\alpha(K \cap J)}{|J|^\alpha} : J \text{ is an interval centered in } K \text{ with } J \subset [0,1]\} = 1.$$

Since $\lambda = \mathcal{P}^\alpha|_K / \mathcal{P}^\alpha(K)$, (1.10) follows immediately from the above equation. □

As mentioned earlier, a special case of Corollary 1.4 is directly proved in earlier work by Feng [6] and the result of Theorem 1.2 is also implicit in that case. In that case the linear Cantor set K satisfies the open set condition with the open set $(0,1)$ which is a special case of the generalized finite condition with respect to the invariant open set $(0,1)$. In another direction, Olsen [19] and the author [20] generalized Feng's works to general self-similar sets in $\mathbb{R}^d$ satisfying the strong separation condition or the open set condition. Similar to the Hausdorff measure case, with suitable modifications if necessary, we may generalize Theorem 1.2 and Corollary 1.4 in the analogous way to general self-similar sets in $\mathbb{R}^d$ satisfying the generalized finite type condition with irreducible weighted incidence matrix. Due to the same reason as that for the Hausdorff measure case, we will not pursue such generalizations here.

4 The maximal and minimal densities

In this section we deal with the exact computation of the Hausdorff measure and the packing measure for a special kind of linear Cantor sets. Let $S_j(x) = \rho_j x + b_j, j =$

$1, \cdots, m$, be a linear contractive IFS on the line satisfying the generalized finite type condition with respect to the open set $\mathcal{O} = (0,1)$ with an irreducible weighted incidence matrix A_α. As before, we write K as its invariant set and use α to denote the common value of the Hausdorff and packing dimensions of K. As stated in [1, 6], We do not allow negative ρ's, i.e., we assume $0 < \rho_j < 1$ for each $j = 1, \cdots, m$. Moreover, without loss of generality and for convenience we assume the images $S_j([0,1])$ are in increasing order, with $S_1(0) = 0$ and $S_m(1) = 1$. To avoid triviality we always assume $m \geq 2$ and $\alpha < 1$.

Define

$$d_{\max} := \sup\{d(J) : J \text{ is an interval with } J \subset [0,1]\}$$

the maximal density of intervals contained in $[0,1]$ and

$$d_{\min} := \inf\{d(J) : J \text{ is an interval centered in } K \text{ with } J \subset [0,1]\}$$

the minimal density of intervals centered in K and contained in $[0,1]$ where $d(J) := \lambda(J)/|J|^\alpha$ denotes the density of the interval J with respect to λ, the α-dimensional normalized Hausdorff measure restricted to K. Then by Corollary 1.3 and Corollary 1.4,

$$\mathcal{H}^\alpha(K) = d_{\max}^{-1} \text{ and } \mathcal{P}^\alpha(K) = d_{\min}^{-1}.$$

Hence our main purpose in this section is to determine the constants $d_{\max}$ and $d_{\min}$.

In this section, we will frequently use the notation $\mathcal{F}_k$ which is the finite field generated by the set $\mathcal{F}_k^0$ for each $k \geq 0$. We call $\mathcal{F}_k$ the *k-th generation field.* For each $k \geq 0$, let $\beta_1^{(k)}, \cdots, \beta_{l_k}^{(k)}$ denote the lengths of the k-th generation islands in increasing order where l_k denotes the cardinality of $\mathcal{F}_k^0$. Let $\beta_{\max}^{(k)} = \max\{\beta_1^{(k)}, \cdots, \beta_{l_k}^{(k)}\}$ and $\beta_{\min}^{(k)} = \min\{\beta_1^{(k)}, \cdots, \beta_{l_k}^{(k)}\}$ be the maximal and minimal length of the k-th generation islands respectively. We write the lengths of the lakes separating the k-th generation islands as $\gamma_1^{(k)}, \cdots, \gamma_{l_k-1}^{(k)}$ in increasing order. Let $\gamma_{\min}^{(k)} = \min\{\gamma_1^{(k)}, \cdots, \gamma_{l_k-1}^{(k)}\}$ be the minimal length of the lakes in $\mathcal{F}_k$. Note that we allow $\gamma_{\min}^{(k)} = 0$ in the case of touching islands, and indeed this case leads to some of complicated and interesting phenomena. Denote by $\gamma_{\min}^{*(k)}$ the minimal length of non-empty lakes in $\mathcal{F}_k$. The identity

$$\beta_1^{(k)} + \cdots + \beta_{l_k}^{(k)} + \gamma_1^{(k)} + \cdots + \gamma_{l_k-1}^{(k)} = 1,$$

the positivity of β's and the non-negativity of γ's are the only restrictions on these parameters. It should be mentioned here that $l_k \to \infty$, $\beta_{\max}^{(k)} \to 0$, $\beta_{\min}^{(k)} \to 0$, $\gamma_{\min}^{(k)} \to 0$ and $\gamma_{\min}^{*(k)} \to 0$ as $k \to \infty$.

The α-dimensional normalized Hausdorff measure λ will play a key role in this section. In view of Remark 3.4, the definition of λ enables us to compute the measure of each island and furthermore the density of each interval in $\mathcal{F}_k$ as an elementary function on parameters of the IFS of K. Moreover, there is an obvious algorithm for finding the maximal or minimal density of intervals in $\mathcal{F}_k$.

By the assumption that K has finite types of islands, there exists a smallest non-negative integer k_0 such that none of the islands in $\mathcal{F}^0_{k_0+1}$ is of a new overlap type. We have the following *blow-up principle* for the density of each interval $J \subset [0,1]$.

Lemma 4.1. *If J is any interval, there exists another interval J', not contained in a (k_0+1)-th generation island, with the same density.*

Proof. If J is contained in a (k_0+1)-th generation island I, there exists a larger island $I' \in \bigcup_{0\le k\le k_0} \mathcal{F}^0_k$ with the same type as I, i.e., $[I'] = [I]$. Let τ be the linear function which maps I onto I', keeping the orientation. Then obviously the image $J' := \tau(J)$ has the same density as J. We iterate this procedure until we obtain J' not lying in any (k_0+1)-th generation island. □

Thus we only need to consider intervals not contained in (k_0+1)-th generation islands.

In order to get uniform results for computing $\mathcal{H}^\alpha(K)$ and $\mathcal{P}^\alpha(K)$, we need the following two additional technical assumptions. We will show that these assumptions are as general as to be satisfied by all the examples illustrated in Section 2.

Assumption A. For each island $I \in \mathcal{F}^0$ and for any two different constitutive intervals $\overline{\mathcal{O}}_{\mathbf{v}}$ and $\overline{\mathcal{O}}_{\mathbf{u}}$ of I, we have $\overline{\mathcal{O}}_{\mathbf{v}} \not\subseteq \overline{\mathcal{O}}_{\mathbf{u}}$ and $\overline{\mathcal{O}}_{\mathbf{u}} \not\subseteq \overline{\mathcal{O}}_{\mathbf{v}}$.

Assumption B. $S_1([0,1]) \cap K = S_1K$ and $S_m([0,1]) \cap K = S_mK$.

Before we give some remarks on these assumptions, we first look at the examples introduced in Section 2.

Example 4.2. If $\{S_j\}_{j=1}^m$ satisfies the open set condition as showed in Example 2.5, Assumption A and Assumption B are naturally satisfied.

The Proof is trivial.

Example 4.3. The IFS $\{S_j\}_{j=1}^3$ on $\mathbb{R}$ defined in Example 2.6 satisfies the condition of Assumption A and Assumption B.

Proof. Assumption A and the second equality of Assumption B hold obviously. Hence we only need to prove the first equality of Assumption B. By the fact $S_1([0,1]) \cap S_3K = \emptyset$, $S_1([0,1]) \cap S_{23}K = \emptyset$ and $S_{13} = S_{21}$, we have

$$\begin{aligned} S_1([0,1]) \cap K &= (S_1([0,1]) \cap S_1K) \cup (S_1([0,1]) \cap S_2K) \\ &= (S_1([0,1] \cap K)) \cup (S_1([0,1]) \cap S_{13}K) \cup (S_1([0,1]) \cap S_2^2K) \\ &= (S_1([0,1] \cap K)) \cup (S_1([0,1] \cap S_3K)) \cup (S_1([0,1]) \cap S_2^2K). \end{aligned}$$

For $n \ge 2$, iterate the above proceedure $n-1$ times, we get

$$\begin{aligned} S_1([0,1]) \cap K = {} & (S_1([0,1] \cap K)) \cup (S_1([0,1] \cap S_3K)) \cup \\ & \cdots \cup (S_1([0,1] \cap S_3^nK)) \cup (S_1([0,1]) \cap S_2^{n+1}K). \end{aligned}$$

Hence we eventually prove that $S_1([0,1])\cap K \subset S_1K$ by the arbitrariness of n. The other direction is obvious. $\Box$

Example 4.4. The IFS $\{S_j\}_{j=1}^3$ on $\mathbb{R}$ defined in Example 2.8 satisfies the condition of Assumption A and Assumption B.

Proof. Assumption A and the second equality of Assumption B hold obviously. Hence we only need to prove the first equality of Assumption B. By the fact $S_1([0,1])\cap S_3K=\emptyset$, $S_1([0,1])\cap S_{23}K=\emptyset$ and $S_{133}=S_{21}$, we have

$$\begin{aligned} S_1([0,1])\cap K &= (S_1([0,1])\cap S_1K)\cup(S_1([0,1])\cap S_2K) \\ &= (S_1([0,1]\cap K))\cup(S_1([0,1])\cap S_{133}K)\cup(S_1([0,1])\cap S_2^2K) \\ &= (S_1([0,1]\cap K))\cup(S_1([0,1]\cap S_3^2K))\cup(S_1([0,1])\cap S_2^2K). \end{aligned}$$

For $n\geq 2$, iterate the above proceedure $n-1$ times, we get

$$\begin{aligned} S_1([0,1])\cap K &= (S_1([0,1]\cap K))\cup(S_1([0,1]\cap S_3^2K))\cup \\ &\qquad \cdots\cup(S_1([0,1]\cap S_3^{2n}K))\cup(S_1([0,1])\cap S_2^{n+1}K). \end{aligned}$$

Hence we eventually prove that $S_1([0,1])\cap K \subset S_1K$ by the arbitrariness of n. The other direction is obvious. $\Box$

Example 4.5. The IFS $\{S_j\}_{j=1}^4$ on $\mathbb{R}$ defined in Example 2.9 satisfies the condition of Assumption A and Assumption B.

Proof. Assumption A holds obviously. For Assumption B, it is also obvious since $S_1([0,1])$ and $S_4([0,1])$ are two first generation islands lying on the left side and right side of $[0,1]$ respectively. $\Box$

Remark 4.6. Under Assumption B, for each positive integer k, we have $S_1^k([0,1])\cap K=S_1^kK$; similarly, for each positive integer k', we have $S_m^{k'}([0,1])\cap K=S_m^{k'}K$.

Proof. By induction, we only need to prove $S_1^2([0,1])\cap K=S_1^2K$. Since $S_1^2([0,1])\cap K\subset S_1([0,1])\cap K=S_1K$, we have $S_1^2([0,1])\cap K=S_1^2([0,1])\cap S_1K=S_1(S_1([0,1])\cap K)=S_1^2K$. The second case can be proved in a similar way. $\Box$

Remark 4.7. For each island $I=[c,d]\in\mathcal{F}^0$, notice that $I=\bigcup_{\mathbf{v}\in V(I)}\overline{\mathcal{O}}_{\mathbf{v}}$. By Assumption A, there is a unique vertex $\mathbf{v}_0\in V(I)$ such that $\overline{\mathcal{O}}_{\mathbf{v}_0}$ has the same left endpoint as I, i.e., $S_{\mathbf{v}_0}(0)=c$. If $V(I)\setminus\{\mathbf{v}_0\}\neq\emptyset$, let $c'=\min\{S_{\mathbf{v}}(0):\mathbf{v}\in V(I)\setminus\{\mathbf{v}_0\}\}$. Obviously $c<c'<d$. Let k be the smallest positive integer such that $S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0,1])\subset[c,c')$ where $\mathbf{i}_0=(1,\cdots,1)$ with $|\mathbf{i}_0|=k$ (This must be done since $S_{\mathbf{v}_0}S_1^k(0)$ is always equal to c). For this $\mathbf{i}_0$, by Remark 4.6, we have

$$S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0,1])\cap K=S_{\mathbf{v}_0}S_{\mathbf{i}_0}K.$$

This can be verified since

$$\begin{aligned} S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0,1])\cap K &= S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0,1])\cap S_{\mathbf{v}_0}K \\ &= S_{\mathbf{v}_0}(S_{\mathbf{i}_0}([0,1])\cap K) \\ &= S_{\mathbf{v}_0}S_{\mathbf{i}_0}K. \end{aligned}$$

Otherwise, if $V(I)\setminus\{\mathbf{v}_0\}=\emptyset$, i.e., $V(I)=\{\mathbf{v}_0\}$, we define $\mathbf{i}_0=\emptyset$. Also, we have $S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0,1])\cap K=S_{\mathbf{v}_0}S_{\mathbf{i}_0}K$.

Similarly, In an analogous way, there is also a unique vertex $\mathbf{v}_1\in V(I)$ such that $\overline{\mathcal{O}}_{\mathbf{v}_1}$ has the same right endpoint as I, i.e., $S_{\mathbf{v}_1}(1)=d$. By a similar discussion, one can find that there also exists a smallest non-negative integer k' such that

$$S_{\mathbf{v}_1}S_{\mathbf{i}_1}([0,1])\cap K=S_{\mathbf{v}_1}S_{\mathbf{i}_1}K$$

where $\mathbf{i}_1=(m,\cdots,m)$ with the length $|\mathbf{i}_1|=k'$.

Remark 4.8. We should mention that in Remark 4.7 the ratios $|S_{\mathbf{v}_0}([0,1])|/|I|$, $|S_{\mathbf{v}_1}([0,1])|/|I|$ and the indices $\mathbf{i}_0,\mathbf{i}_1$ are dependent merely on the overlap type of I and are independent on the choice of I. In order to emphasize the relation between $\mathbf{v}_0,\mathbf{v}_1,\mathbf{i}_0,\mathbf{i}_1$ and I, we replace $\mathbf{v}_0,\mathbf{v}_1,\mathbf{i}_0,\mathbf{i}_1$ by $\mathbf{v}_0(I),\mathbf{v}_1(I),\mathbf{i}_0(I),\mathbf{i}_1(I)$ respectively. Since the number of the overlap types is finite, we could define two positive numbers $\eta_1\le 1$ and $\eta_2\le 1$ as follows.

$$\eta_1:=\min_{I\in\mathcal{F}^0}\{\frac{\rho_{\mathbf{v}_0(I)}}{|I|},\frac{\rho_{\mathbf{v}_1(I)}}{|I|}\}>0,$$

and

$$\eta_2:=\min_{I\in\mathcal{F}^0}\{\rho_{\mathbf{i}_0(I)},\rho_{\mathbf{i}_1(I)}\}>0.$$

Write $\eta:=\eta_1\eta_2$ which will be used later. Here $0<\eta\le 1$.

In the following, we will always assume Assumption A and Assumption B.

Under these assumptions, we then have the following type of blow-up principle.

Lemma 4.9 *If $J\subset[0,1]$ is any interval of the form $[0,x]$, there exists another interval $J'=[0,x']$ of the same form with $\rho_1<x'\le 1$ such that $d(J')=d(J)$; similarly, if J is any interval of the form $[y,1]$, there exists another interval $J'=[y',1]$ of the same form with $0\le y'<1-\rho_m$ such that $d(J')=d(J)$.*

Proof. By Assumption B, if $J\subset S_1([0,1])$, $S_1^{-1}J$ is a larger interval of the same density. We iterate this procedure until we obtain J' not lying in $S_1([0,1])$. The proof of the second case is similar. □

For an island $I=[c,d]\subset[0,1]$, we introduce the following notations.

$$\underline{D}_0(I)=\inf_{0<x\le 1}\{d([c,c+x(d-c)])\}\text{ and }\underline{D}_1(I)=\inf_{0<x\le 1}\{d([d-x(d-c),d])\}.$$

Obviously, if I_1 and I_2 are two islands with the same overlap type, i.e., $[I_1] = [I_2]$,

$$\underline{D}_0(I_1) = \underline{D}_0(I_2) \text{ and } \underline{D}_1(I_1) = \underline{D}_1(I_2).$$

Hence the notations $\underline{D}_0(I)$ and $\underline{D}_1(I)$ depend only on the overlap type of the island I. Thus we could define the following constants. For $1 \leq i \leq q$, define

$$\underline{D}_0^i := \underline{D}_0(I) \text{ and } \underline{D}_1^i := \underline{D}_1(I)$$

where I is an arbitrary $\mathcal{T}_i$ type island. Here these notations are independent of the choice of the island I. The following lemma shows Assumption A and Assumption B will ensure that all $\underline{D}_0^i$'s are equal and all $\underline{D}_1^i$'s are also equal.

Lemma 4.10.

$$\underline{D}_0^1 = \cdots = \underline{D}_0^q \text{ and } \underline{D}_1^1 = \cdots = \underline{D}_1^q. \tag{4.1}$$

Proof. Fix $2 \leq i \leq q$. We prove $\underline{D}_0^1 = \underline{D}_0^i$. Take a $\mathcal{T}_i$ type island $I := [c, d]$. Now we turn to prove $\underline{D}_0([0,1]) = \underline{D}_0(I)$ since $[0, 1]$ is of $\mathcal{T}_1$ type.

Using Remark 4.7, we have $S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0, 1]) \cap K = S_{\mathbf{v}_0}S_{\mathbf{i}_0}K$ where $\mathbf{v}_0 = \mathbf{v}_0(I)$ and $\mathbf{i}_0 = \mathbf{i}_0(I)$ are the same notations as those discussed in Remark 4.7 and Remark 4.8. Hence for any $0 < x \leq 1$, since $S_{\mathbf{v}_0}S_{\mathbf{i}_0}([0, 1]) \cap K$ is similar to K, we have

$$d([0, x]) = d([c, c + x(S_{\mathbf{v}_0}S_{\mathbf{i}_0}(1) - c)]),$$

which yields that $\underline{D}_0^i \leq \underline{D}_0^1$ by the arbitrariness of x and the fact that $[c, c+x(S_{\mathbf{v}_0}S_{\mathbf{i}_0}(1) - c)] \subset I$.

On the other hand, since $I = \bigcup_{\mathbf{v}\in V(I)} S_{\mathbf{v}}([0, 1])$, we denote by $c = a_1 < a_2 < \cdots < a_n$ the left endpoints of all constitutive intervals of I in increasing order. (By Assumption A, it is impossible that some distinct constitutive intervals share a common left endpoint.) For any interval $[c, z] \subset I$, choose a largest a_i such that $a_i \leq z$. If $a_i < z$,

$$\begin{aligned} d([c, z]) &= \frac{\lambda([a_1, a_2]) + \cdots + \lambda([a_{i-1}, a_i]) + \lambda([a_i, z])}{((a_2 - a_1) + \cdots + (a_i - a_{i-1}) + (z - a_i))^\alpha} \\ &\geq \frac{\lambda([a_1, a_2]) + \cdots + \lambda([a_{i-1}, a_i]) + \lambda([a_i, z])}{(a_2 - a_1)^\alpha + \cdots + (a_i - a_{i-1})^\alpha + (z - a_i)^\alpha} \\ &\geq \min\{d([a_1, a_2]), \cdots, d([a_{i-1}, a_i]), d([a_i, z])\} \\ &\geq \underline{D}_0^1. \end{aligned}$$

The last inequality follows from the fact that for each $\mathbf{v} \in V(I)$, $S_{\mathbf{v}}K \subset S_{\mathbf{v}}([0, 1]) \cap K$ and $S_{\mathbf{v}}K$ is similar to K. In the case $a_i = z$, we have the same result by a similar discussion. The arbitrariness of z yields that $\underline{D}_0^i \geq \underline{D}_0^1$.

Hence we have $\underline{D}_0^1 = \underline{D}_0^i$. By the arbitrariness of $2 \le i \le q$, we get the first equality in (4.1). The second equality can be proved in a similar way. $\square$

Thus we could define the common value of $\underline{D}_0^1, \cdots, \underline{D}_0^q$ as $\underline{D}_0$ and the common value of $\underline{D}_1^1, \cdots, \underline{D}_1^q$ as $\underline{D}_1$. We would like to characterize $\underline{D}_0$ and $\underline{D}_1$ by the parameters β's and γ's.

Lemma 4.11. *Let k be the smallest integer such that $\beta_1^{(k)} \le \rho_1$. Then*

$$\underline{D}_0 = \min\{d([0,x]) : x > 0 \text{ and } [0,x] \in \mathcal{F}_k\}.$$

Similarly, let k' be the smallest integer such that $\beta_{l_{k'}}^{(k')} \le \rho_m$. Then

$$\underline{D}_1 = \min\{d([y,1]) : y < 1 \text{ and } [y,1] \in \mathcal{F}_{k'}\}.$$

Proof. For simplicity, we only prove the first equality. By the definition of $\underline{D}_0$, we note that

$$\underline{D}_0 = \inf_{0<x\le 1} d([0,x]).$$

By Assumption B and the blow-up principle Lemma 4.9, we only need to consider the interval $[0,x]$ with $\rho_1 < x \le 1$. Since $d([0,x])$ is a continuous function of x, $d([0,x])$ attains its minimum $\underline{D}_0$ at some $x_0 \in [\rho_1, 1]$. Furthermore noting that $d([0,1]) = \underline{D}_0$ whenever $d([0,\rho_1]) = \underline{D}_0$, we can assume $x_0 > \rho_1$.

It is clear that the point x_0 can not fall in a non-empty lake of $\mathcal{F}_k$ because then $[0,x_0]$ would not have minimal density. Therefore there exists a k-th generation island $I = [c,d]$ such that $x_0 \in [c,d]$. Here $c > 0$ because otherwise $x_0 \le \beta_1^{(k)} \le \rho_1$ which contradicts $x_0 > \rho_1$. Take $u = x_0 - c$. Assume that $u > 0$. Then

$$\begin{aligned} d([0,x_0]) &= \frac{\lambda([0,c]) + \lambda([c,x_0])}{(c+u)^\alpha} \\ &> \frac{\lambda([0,c]) + \lambda([c,x_0])}{c^\alpha + u^\alpha} \\ &\ge \min\{\frac{\lambda([0,c])}{c^\alpha}, \frac{\lambda([c,x_0])}{u^\alpha}\} \\ &= \min\{d([0,c]), d([c,x_0])\} \\ &\ge \underline{D}_0, \end{aligned}$$

which contradicts the minimality of $d([0,x_0])$. Hence $u = 0$. Thus

$$\underline{D}_0 = \min\{d([0,x]) : x > 0 \text{ and } [0,x] \in \mathcal{F}_k\}. \quad \square$$

We also need to consider the maximal value of the density $d(J)$ whenever J is of the form $[0,x]$ or $[y,1]$. For this purpose, define

$$\overline{D}_0 = \sup\{d([0,x]) : 0 < x \le 1\} \text{ and } \overline{D}_1 = \sup\{d([y,1]) : 0 \le y < 1\}.$$

We would also like to characterize $\overline{D}_0$ and $\overline{D}_1$ by the parameters β's and γ's. The following elementary calculus lemma by Ayer & Strichartz is useful.

Lemma 4.12[1]**.** *Suppose* $0 < \alpha < 1$, $p \leq p_0$, $a \geq a_0$, $\kappa > 0$ *and* $y \geq \kappa x^\alpha$. *Then*

$$0 < x \leq (\frac{a_0 \kappa}{p_0})^{\frac{1}{1-\alpha}} \tag{4.2}$$

implies

$$\frac{p-y}{(a-x)^\alpha} < \frac{p}{a^\alpha}. \tag{4.3}$$

To make this paper self-contained, we give the proof of Lemma 4.12 as follows.

Proof of Lemma 4.12. Consider the function $f(x) = (p - \kappa x^\alpha)/(a-x)^\alpha$. Noting that $(p-y)/(a-x)^\alpha \leq f(x)$ by the assumption $y \geq \kappa x^\alpha$. And by $p/a^\alpha = f(0)$, it suffices to show $f'(x) < 0$ on the interval $0 < x < (a_0\kappa/p_0)^{1/(1-\alpha)}$. This can be verified by a direct computation. □

Using the above lemma, we have the following result concerning $\overline{D}_0$ and $\overline{D}_1$.

Lemma 4.13. *Let k be the smallest integer such that* $\beta^{(k)}_{\max} \leq (\rho_1 \underline{D}_1)^{1/(1-\alpha)}$. *Then*

$$\overline{D}_0 = \max\{d([0,x]) : x > 0 \textit{ and } [0,x] \in \mathcal{F}_k\}.$$

Similarly, let k' be the smallest integer such that $\beta^{(k')}_{\max} \leq (\rho_m \underline{D}_0)^{1/(1-\alpha)}$. *Then*

$$\overline{D}_1 = \max\{d([y,1]) : y < 1 \textit{ and } [y,1] \in \mathcal{F}_{k'}\}.$$

Proof. For simplicity, we only prove the first equality. By the blow-up principle Lemma 4.9 we can take $x_0 > \rho_1$ such that the interval $[0, x_0]$ has the maximal density (By compactness of $[\rho_1, 1]$, the maximum is attained). The point x_0 cannot fall in a non-empty lake of $\mathcal{F}_k$ because, if so, $[0, x_0]$ would not have maximal density. Let $[0, a]$ be the smallest interval in $\mathcal{F}_k$ that contains $[0, x_0]$. Then $x_0 = a - x$ for some $x \leq \beta^{(k)}_{\max}$ because $\beta^{(k)}_{\max}$ is the length of the largest island in $\mathcal{F}^0_k$. Set $p = \lambda([0,a])$ and $y = \lambda([a-x, a])$, then $d([0, a-x]) = (p-y)/(a-x)^\alpha$ and $d([0,a]) = p/a^\alpha$. Thus the conclusion (4.3) of Lemma 4.12 would give $d([0,a-x]) < d([0,a])$ unless $x = 0$, which implies that $[0,a]$ attains the maximal density.

To complete the proof we will verify the hypothesis of Lemma 4.12 with $p_0 = 1$, $a_0 = \rho_1$ and $\kappa = \underline{D}_1$. We already know $a \geq \rho_1$, and $p \leq 1$ is trivial. To verify $y \geq \kappa x^\alpha$, we observe that $y/x^\alpha = d([a-x,a])$, and by the definition of $\underline{D}_1$, we immediately get $d([a-x,a]) \geq \underline{D}_1$. The hypothesis of Lemma 4.12 is verified, and condition (4.2) follows from $x \leq \beta^k_{\max}$ and the hypothesis $\beta^{(k)}_{\max} \leq (\rho_1 \underline{D}_1)^{1/(1-\alpha)}$. □

4.1 The maximal density and the Hausdorff measure

For $k \geq 0$, notice that $\gamma_{\min}^{(k)}$ is the minimal value of lengths of the lakes separating the k-th generation islands. If $\gamma_{\min}^{(k)} = 0$, there must exist touching k-th generation islands. And this will make things complicated. As stated before, we denote by k_0 the smallest non-negative integer such that none of the islands in $\mathcal{F}_{k_0+1}^0$ are of new overlap type. First, in the case that all lakes are non-zero, i.e., $\gamma_{\min}^{(k_0+1)} > 0$, we have the following result.

Theorem 4.14. *Assume* $\gamma_{\min}^{(k_0+1)} > 0$*, and let* $k \geq k_0 + 1$ *be the smallest integer such that*

$$2\beta_{\max}^{(k)} \leq (\gamma_{\min}^{(k_0+1)} \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}}. \tag{4.4}$$

Then the maximal density $d_{\max}$ *is attained for an interval in* $\mathcal{F}_k$*.*

Proof. By the blow-up principle Lemma 4.1 we may focus attention to intervals containing at least one lake of $\mathcal{F}_{k_0+1}$. So we have the lower bound $\gamma_{\min}^{(k_0+1)}$ for the length of the interval, which implies by compactness that the maximal density $d_{\max}$ is attained. If $[x_1, x_2]$ is an interval of maximal density, we let $[z_1, z_2]$ be the smallest interval in $\mathcal{F}_k$ containing $[x_1, x_2]$. Write $a = z_2 - z_1$ for the length of the interval, $x = (z_2 - z_1) - (x_2 - x_1)$ for the difference of the lengths, $p = \lambda([z_1, z_2])$ and $y = \lambda([z_1, x_1]) + \lambda([x_2, z_2])$, so that $d([x_1, x_2]) = (p - y)/(a - x)^\alpha$ and $d([z_1, z_2]) = p/a^\alpha$. By an analogous way in proving Lemma 4.13, we will complete the proof by applying Lemma 4.12.

We take $p_0 = 1$ and $a_0 = \gamma_{\min}^{(k_0+1)}$, so that $a \geq a_0$ and $p \leq p_0$. We choose $\kappa = \min\{\underline{D}_0, \underline{D}_1\}$. For the right side interval $[x_2, z_2]$ we have

$$\frac{\lambda([x_2, z_2])}{(z_2 - x_2)^\alpha} \geq \underline{D}_1,$$

and similarly for the left side interval $[z_1, x_1]$ we have

$$\frac{\lambda([z_1, x_1])}{(x_1 - z_1)^\alpha} \geq \underline{D}_0.$$

Thus we have

$$y \geq \min\{\underline{D}_0, \underline{D}_1\}((x_1 - z_1)^\alpha + (z_2 - x_2)^\alpha) \geq \kappa x^\alpha$$

since $0 < \alpha < 1$. Thus the hypothesis of Lemma 4.12 is verified, and condition (4.2) follows from (4.4) since x is the sum of two terms, $x_1 - z_1$ and $z_2 - x_2$, each being at most $\beta_{\max}^{(k)}$. □

Now we turn to discuss the case that there exist touching islands. First we can still obtain a result if we assume a logarithmic arithmetic relation between ρ_1 and ρ_m. In the following, η is the positive number defined in Remark 4.8.

Theorem 4.15. *Suppose there exist positive integers* n_1 *and* n_m *such that* $\rho_1^{n_1} = \rho_m^{n_m}$*. Let* k *be the smallest integer such that*

$$2\beta_{\max}^{(k)} \leq (\eta \beta_{\min}^{(k_0+1)} \rho_1^{n_1} \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}}.$$

Then the maximal density $d_{\max}$ *is attained for an interval in* $\mathcal{F}_k$*.*

Proof. We claim that it suffices to consider intervals of length at least $\eta \beta_{\min}^{(k_0+1)} \rho_1^{n_1}$. To see this we need a variant of the blow-up principle that shows how to replace smaller intervals with larger intervals of greater density.

Start with any interval I_0 not contained in a (k_0+1)-th generation island. If it actually contains a (k_0+1)-th generation island, its length is at least $\beta_{\min}^{(k_0+1)}$, and we are done. If not, it begins at a point in I and ends at a point in I' where I, I' are two adjacent (k_0+1)-th generation islands. Denote by L the lake separating I and I'. Using Remark 4.7, there exists a vertex $\mathbf{v} \in V(I)$ and an index $\mathbf{i} = (m, \cdots, m)$ such that $S_{\mathbf{v}}([0,1]) \subset I$ has the same right endpoint as that of I and $S_{\mathbf{v}}S_{\mathbf{i}}([0,1]) \cap K = S_{\mathbf{v}}S_{\mathbf{i}}K$. Similarly, Remark 4.7 also ensures that there exists a vertex $\mathbf{v}' \in V(I')$ and an index $\mathbf{i}' = (1, \cdots, 1)$ such that $S_{\mathbf{v}'}([0,1]) \subset I'$ has the same left endpoint as that of I' and $S_{\mathbf{v}'}S_{\mathbf{i}'}([0,1]) \cap K = S_{\mathbf{v}'}S_{\mathbf{i}'}K$. For simplicity, we denote $S_{\mathbf{v}}S_{\mathbf{i}}([0,1])$ and $S_{\mathbf{v}'}S_{\mathbf{i}'}([0,1])$ by $\widetilde{I}$ and $\widetilde{I}'$ respectively. (These notations will be used again in Lemma 4.17. and Theorem 4.18.) Now consider the intervals $J = S_{\mathbf{v}}S_{\mathbf{i}}S_m^{n_m}([0,1])$ and $J' = S_{\mathbf{v}'}S_{\mathbf{i}'}S_1^{n_1}([0,1])$ which lie on the extreme ends of the lake L separating $\widetilde{I}$ and $\widetilde{I}'$. These intervals have length $\rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_m^{n_m}$ and $\rho_{\mathbf{v}'}\rho_{\mathbf{i}'}\rho_1^{n_1}$. Moreover, $J \cap K$ and $J' \cap K$ are similar to K. If our interval I_0 contains one of them, we are done.

Next suppose our interval begins with a point in J and ends with a point in J', say $I_0 = J_0 \cup L \cup J_0'$ where $J_0 = I_0 \cap J$ and $J_0' = I_0 \cap J'$ and L is the lake separating J and J'. We generate another interval $I_1 = J_1 \cup L \cup J_1'$ by blowing up J_0 to J_1 and J_0' to J_1' by a factor $\rho_1^{-n_1} = \rho_m^{-n_m}$. Specifically, we set $J_1 = S_{\mathbf{v}}S_{\mathbf{i}}S_m^{-n_m}(S_{\mathbf{v}}S_{\mathbf{i}})^{-1}J_0$ and $J_1' = S_{\mathbf{v}'}S_{\mathbf{i}'}S_1^{-n_1}(S_{\mathbf{v}'}S_{\mathbf{i}'})^{-1}J_0'$. Note that $S_{\mathbf{v}}S_{\mathbf{i}}S_m^{-n_m}(S_{\mathbf{v}}S_{\mathbf{i}})^{-1}$ maps J onto $\widetilde{I}$ and fixes the right endpoint, while $S_{\mathbf{v}'}S_{\mathbf{i}'}S_1^{-n_1}(S_{\mathbf{v}'}S_{\mathbf{i}'})^{-1}$ maps J' onto $\widetilde{I}'$ and fixes the left endpoint. So I_1 is an interval. We have

$$d(I_0) = \frac{\lambda(J_0) + \lambda(J_0')}{(|J_0| + |L| + |J_0'|)^{\alpha}},$$

while

$$\begin{aligned} d(I_1) &= \frac{\rho_m^{-n_m\alpha}\lambda(J_0) + \rho_1^{-n_1\alpha}\lambda(J_0')}{(\rho_m^{-n_m}|J_0| + |L| + \rho_1^{-n_1}|J_0'|)^{\alpha}} \\ &= \frac{\lambda(J_0) + \lambda(J_0')}{(|J_0| + \rho_1^{n_1}|L| + |J_0'|)^{\alpha}}. \end{aligned}$$

So $d(I_1) \geq d(I_0)$. By iterating this blow-up construction we eventually arrive at an interval containing either J or J' whose density is greater than the original interval I_0.

Hence we only need to consider intervals containing either J or J' which have length at least $\min\{\rho_{\mathbf{v}}\rho_{\mathbf{i}}, \rho_{\mathbf{v}'}\rho_{\mathbf{i}'}\}\rho_1^{n_1}$. Since by the definition of the constant η, $\min\{\rho_{\mathbf{v}}\rho_{\mathbf{i}}, \rho_{\mathbf{v}'}\rho_{\mathbf{i}'}\}$ has the lower bound $\eta\beta_{\min}^{(k_0+1)}$. This completes the proof that it suffices to look at intervals of length at least $\eta\beta_{\min}^{(k_0+1)}\rho_1^{n_1}$.

The rest of the argument is identical to the proof of Theorem 4.14, except that we take $a_0 = \eta\beta_{\min}^{(k_0+1)}\rho_1^{n_1}$. $\square$

We consider now the case when the contraction ratios ρ_1 and ρ_m do not satisfy the arithmetic condition of Theorem 4.15. Another elementary calculus lemma proved by Ayer & Strichartz in [1] will be needed.

Lemma 4.16[**1**]**.** *Let $a, a', q, q' > 0$, and $0 < \alpha < 1$. $F(x)$ is a function*

$$F(x) = \frac{q + q'x^{\alpha}}{(a + a'x)^{\alpha}} \tag{4.5}$$

of positive variables. Then F attains the maximal value of

$$((\frac{q}{a^{\alpha}})^{\frac{1}{1-\alpha}} + (\frac{q'}{a'^{\alpha}})^{\frac{1}{1-\alpha}})^{1-\alpha}$$

at the point

$$x_0 = (\frac{aq'}{a'q})^{\frac{1}{1-\alpha}}.$$

Furthermore, $F(x)$ is strictly increasing on $0 \leq x \leq x_0$ and strictly decreasing on $x > x_0$.

Proof. This can be done by a directly computation of $F'(x)$. $\square$

Lemma 4.17. *Suppose I and I' are two (k_0+1)-th generation touching islands, and ρ_1 and ρ_m are non-arithmetic in the sense that $\rho_1^{n_1} \neq \rho_m^{n_m}$ for any positive integers n_1 and n_m. $\widetilde{I}$ and $\widetilde{I}'$ are the same as that defined in the proof of Theorem 4.15. Then the maximal density of intervals beginning in $\widetilde{I}$ and ending in $\widetilde{I}'$ is*

$$(\overline{D}_0^{\frac{1}{1-\alpha}} + \overline{D}_1^{\frac{1}{1-\alpha}})^{1-\alpha}. \tag{4.6}$$

Proof. Any such interval can be written as $I_0 = \widetilde{I}_0 \cup \widetilde{I}_0'$ where $\widetilde{I}_0 \subset \widetilde{I}$ ends at the right endpoint of $\widetilde{I}$, and $\widetilde{I}_0' \subset \widetilde{I}'$ begins at the left endpoint of $\widetilde{I}'$. Let $\mathbf{v}, \mathbf{v}', \mathbf{i}, \mathbf{i}'$ be the notations used in the proof of Theorem 4.15. So $S_{\mathbf{v}}S_{\mathbf{i}}([0,1]) = \widetilde{I}$ and $S_{\mathbf{v}'}S_{\mathbf{i}'}([0,1]) = \widetilde{I}'$. For any positive integers k and k' we can form the interval

$$I(k,k') = S_{\mathbf{v}}S_{\mathbf{i}}S_m^k(S_{\mathbf{v}}S_{\mathbf{i}})^{-1}\widetilde{I}_0 \cup S_{\mathbf{v}'}S_{\mathbf{i}'}S_1^{k'}(S_{\mathbf{v}'}S_{\mathbf{i}'})^{-1}\widetilde{I}_0',$$

which contracts $\widetilde{I}_0$ by a factor of ρ_m^k and $\widetilde{I}_0'$ by a factor of $\rho_1^{k'}$, keeping their common endpoint fixed. Then

$$d(I(k,k')) = \frac{\rho_m^{k\alpha}\lambda(\widetilde{I}_0) + \rho_1^{k'\alpha}\lambda(\widetilde{I}_0')}{(\rho_m^k|\widetilde{I}_0| + \rho_1^{k'}|\widetilde{I}_0'|)^{\alpha}} = \frac{\rho_m^{k\alpha}q + \rho_1^{k'\alpha}q'}{(\rho_m^k a + \rho_1^{k'}a')^{\alpha}}, \tag{4.7}$$

where $a = |\widetilde{I}_0|$, $a' = |\widetilde{I}_0'|$ and $q = \lambda(\widetilde{I}_0)$, $q' = \lambda(\widetilde{I}_0')$. Notice that this is exactly of the form (4.5) with $x = \rho_1^{k'}\rho_m^{-k}$, and by the non-arithmetic hypothesis that x takes on a dense set of values on the positive line. Thus (4.7) has maximal value

$$(d(\widetilde{I}_0)^{\frac{1}{1-\alpha}} + d(\widetilde{I}_0')^{\frac{1}{1-\alpha}})^{1-\alpha} \tag{4.8}$$

by Lemma 4.16.

Since (4.8) is an increasing function of $d(\widetilde{I}_0)$ and $d(\widetilde{I}_0')$, and by the fact that $\widetilde{I} \cap K$ and $\widetilde{I}' \cap K$ are similar to K, it is clear that its maximum is attained when $d(\widetilde{I}_0)$ and $d(\widetilde{I}_0')$ assume their maxima, and these are clearly $\overline{D}_1$ and $\overline{D}_0$. Hence the maximal density of intervals beginning in $\widetilde{I}$ and ending in $\widetilde{I}'$ is $(\overline{D}_0^{1/(1-\alpha)} + \overline{D}_1^{1/(1-\alpha)})^{1-\alpha}$. □

Theorem 4.18. *Suppose $\gamma_{\min}^{(k_0+1)} = 0$, and ρ_1 and ρ_m are non-arithmetic. Let k, k_1 and k_2 be the smallest integers such that*

$$2\beta_{\max}^{(k)} \le (\eta \min\{\beta_{\min}^{(k_0+1)}, \gamma_{\min}^{*(k_0+1)}\} \cdot \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}},$$

$$\beta_{\max}^{(k_1)} \le (\rho_1 \underline{D}_1)^{\frac{1}{1-\alpha}},$$

and

$$\beta_{\max}^{(k_2)} \le (\rho_m \underline{D}_0)^{\frac{1}{1-\alpha}}.$$

Then the maximal density $d_{\max}$ is equal to the maximum of the finite set of values $d(I)$ as I varies over all intervals in $\mathcal{F}_k$, and $(d(I_1)^{1/(1-\alpha)} + d(I_2)^{1/(1-\alpha)})^{1-\alpha}$ as I_1 varies over all intervals of the form $[0, x]$ in $\mathcal{F}_{k_1}$ and I_2 varies over all intervals of the form $[y, 1]$ in $\mathcal{F}_{k_2}$.

Proof. It follows from $\gamma_{\min}^{(k_0+1)} = 0$ that there exist (k_0+1)-th generation touching islands in $\mathcal{F}_{k_0+1}^0$. If I and I' are two such islands with I lying on the left side of I', we apply Lemma 4.17, which means we have to consider the values of (4.6). But by Lemma 4.13, $\overline{D}_0$ is attained for an interval of the form $[0, x]$ in $\mathcal{F}_{k_1}$, and similarly, $\overline{D}_1$ is attained for an interval of the form $[y, 1]$ in $\mathcal{F}_{k_2}$.

For every two touching (k_0+1)-th generation islands I and I', denote $\widetilde{I}$ and $\widetilde{I}'$ the corresponding subsets of I and I' respectively as that discussed in Lemma 4.17. We need to consider all intervals beginning in $I \setminus \widetilde{I}$ and ending in I', or beginning in I and ending in $I' \setminus \widetilde{I}'$ for some touching (k_0+1)-th generation islands I and I', and all intervals that contain either a non-zero lake of $\mathcal{F}_{k_0+1}$ or a (k_0+1)-th generation island. In the first case, intervals have length at least $\min\{|\widetilde{I}|, |\widetilde{I}'|\}$ which always greater than $\beta_{\min}^{(k_0+1)}\eta$ by Remark 4.8. And in the second case, intervals have lengths bounded below by $\min\{\beta_{\min}^{(k_0+1)}, \gamma_{\min}^{*(k_0+1)}\}$. Hence the lengths of all the above two kinds of intervals are bounded below by $\eta \min\{\beta_{\min}^{(k_0+1)}, \gamma_{\min}^{*(k_0+1)}\}$. Then by a slightly variant of the proof of Theorem 4.14, the maximal density over all such intervals is attained by an interval in $\mathcal{F}_k$. □

4.2 The minimal centered density and the packing measure

First, we give a lemma concerning the relation between $d_{\min}$ and $\underline{D}_0, \underline{D}_1$.

Lemma 4.19. $d_{\min} \leq 2^{-\alpha} \min\{\underline{D}_0, \underline{D}_1\}$.

Proof. It suffices to show that there exist J_0, J_1 centered in K, contained in $[0,1]$, such that $d(J_0) = 2^{-\alpha}\underline{D}_0$ and $d(J_1) = 2^{-\alpha}\underline{D}_1$. For simplicity, we only prove the first equality. By Lemma 4.11, let k be the smallest integer such that $\beta_1^{(k)} \leq \rho_1$, then there exists x_0 with $[0, x_0] \subset \mathcal{F}_k$ such that $d([0,x_0]) = \underline{D}_0$. Since $\alpha < 1$, there must exist at least one non-empty lake in $\mathcal{F}_1$, i.e., there exists $1 \leq i \leq l_1 - 1$ with $\gamma_i^{(1)} > 0$, where $\gamma_i^{(1)}$ is the length of the lake separating the i-th and $(i+1)$-th first generation islands I and I'. From the discussion in Remark 4.7, there is a unique vertex $\mathbf{v} \in V(I')$ and an index $\mathbf{i} = (1, \cdots, 1)$ such that the sub-interval $S_{\mathbf{v}}S_{\mathbf{i}}([0,1]) \subset I'$ has the same left endpoint as that of I' and furthermore, $S_{\mathbf{v}}S_{\mathbf{i}}([0,1]) \cap K = S_{\mathbf{v}}S_{\mathbf{i}}K$, i.e., $S_{\mathbf{v}}S_{\mathbf{i}}([0,1]) \cap K$ is similar to K with a contraction ratio $\rho_{\mathbf{v}}\rho_{\mathbf{i}}$. Choose a non-negative integer k' large enough so that $\rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0 < \gamma_i^{(1)}$. Hence the interval $[S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0) - \rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0, S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0)]$ is contained in the lake separating I and I'. Thus $\lambda([S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0) - \rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0, S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0)]) = 0$. Define

$$J_0 := [S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0) - \rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0, S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0) + \rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0].$$

Since the interval $S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}([0,1]) \cap K$ is also similar to K, we have

$$\begin{aligned} d(J_0) &= \frac{\lambda([S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0) - \rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0, S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}(0) + \rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0])}{2^{\alpha}(\rho_{\mathbf{v}}\rho_{\mathbf{i}}\rho_1^{k'}x_0)^{\alpha}} \\ &= \frac{\lambda(S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}([0,x_0]))}{2^{\alpha}|S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}([0,x_0])|^{\alpha}} \\ &= 2^{-\alpha}d(S_{\mathbf{v}}S_{\mathbf{i}}S_1^{k'}([0,x_0])) \\ &= 2^{-\alpha}d([0,x_0]) \\ &= 2^{-\alpha}\underline{D}_0, \end{aligned}$$

which concludes the proof. □

Lemma 4.20. *Let* $J \subset [0,1]$ *be an interval centered in* K *and not contained in any* (k_0+1)*-th generation island. If* $|J| \leq \min\{\beta_{\min}^{(k_0+1)}, \gamma_{\min}^{*(k_0+1)}\}$,

$$d(J) \geq 2^{-\alpha}\min\{\underline{D}_0, \underline{D}_1\}.$$

Proof. For each interval $J \subset [0,1]$ centered in K and not contained in any (k_0+1)-th generation island with $|J| \leq \min\{\beta_{\min}^{(k_0+1)}, \gamma_{\min}^{*(k_0+1)}\}$. It is clear that there are only three possible cases for J.

Case 1: There exist two touching (k_0+1)-th generation islands I_1 and I_2 with I_1 lying on the left side of I_2, and $J = J_1 \cup J_2$, where $J_1 \subset I_1$ and $J_2 \subset I_2$.

In this case, we have

$$d(J) = \frac{\lambda(J_1) + \lambda(J_2)}{(|J_1| + |J_2|)^\alpha} > \frac{\lambda(J_1) + \lambda(J_2)}{|J_1|^\alpha + |J_2|^\alpha} \geq \min\{d(J_1), d(J_2)\} \geq \min\{\underline{D}_0, \underline{D}_1\}$$

by the definition of the constants $\underline{D}_0, \underline{D}_1$. It follows that $d(J) > \min\{\underline{D}_0, \underline{D}_1\}$.

Case 2: There exist two separate (k_0+1)-th generation islands I_1 and I_2 with I_1 lying on the left side of I_2, and $J = J_1 \cup J_2$, where $J_1 \subset I_1$, $J_2 \subset L$, and L is the lake separating I_1 and I_2.

In this case, we have $|J_1| \geq |J_2|$ since J is centered in K. Thus

$$d(J) = \frac{\lambda(J_1)}{(|J_1| + |J_2|)^\alpha} \geq \frac{\lambda(J_1)}{2^\alpha |J_1|^\alpha} = 2^{-\alpha} d(J_1) \geq 2^{-\alpha} \underline{D}_1$$

by the definition of $\underline{D}_1$.

Case 3: There exist two separate (k_0+1)-th generation islands I_1 and I_2 with I_1 lying on the left side of I_2, and $J = J_1 \cup J_2$, where $J_1 \subset L$, $J_2 \subset I_2$, and L is the lake separating I_1 and I_2.

In this case, we have $d(J) \geq 2^{-\alpha}\underline{D}_0$ by a discussion similar to Case 2.

Combining the above discussion completes the proof. □

For convenience, denote all (k_0+1)-th generation islands in increasing order by $I_1, \cdots, I_{l_{k_0+1}}$. For each $1 \leq i \leq l_{k_0+1}$, write $I_i = [a_i, b_i]$.

Lemma 4.21. *Assume* $d_{\min} < 2^{-\alpha} \min\{\underline{D}_0, \underline{D}_1\}$. *Then*

$$d_{\min} = \min_{1 \leq i_1 < i_2 < l_{k_0+1}} \frac{\sum_{i=i_1+1}^{i_2} \lambda([a_i, b_i])}{(a_{i_2+1} - b_{i_1} - 2\,dist(\frac{a_{i_2+1}+b_{i_1}}{2}, K))^\alpha}.$$

Proof. Since $d_{\min} < 2^{-\alpha} \min\{\underline{D}_0, \underline{D}_1\}$, by the blow-up principle Lemma 4.1 and Lemma 4.20, we only need to consider intervals with lengths greater than $\min\{\beta_{\min}^{(k_0+1)}, \gamma_{\min}^{*(k_0+1)}\}$ which are centered in K and not contained in any (k_0+1)-th generation island.

By the compactness of K, there exists a such interval $J_0 = [a_0, b_0]$ such that $d_{\min} = d(J_0)$. First we prove the following statements.

(1) Either $a_0 \in \{b_i : 1 \leq i \leq l_{k_0+1} - 1\}$ or a_0 is contained in one lake;

(2) Either $b_0 \in \{a_i : 2 \leq i \leq l_{k_0+1}\}$ or b_0 is contained in one lake.

For simplicity we only prove (1). The statement (2) will follow by a similar argument. Assume that (1) is not true. Then there exists a $1 \leq i \leq l_{k_0+1} - 1$ such that $a_0 \in [a_i, b_i)$. In the following we will lead to a contradiction. We first claim $(a_0+b_0)/2 > b_i$. Otherwise

$(a_0+b_0)/2 \in [a_i, b_i]$, then

$$d([a_0,b_0]) = \frac{\lambda([a_0,b_0])}{(b_0-a_0)^\alpha} \geq \frac{\lambda([a_0,b_i])}{2^\alpha(b_i-a_0)^\alpha} = 2^{-\alpha} d([a_0,b_i]) \geq 2^{-\alpha}\underline{D}_1,$$

which contradicts the assumption $d_{\min} < 2^{-\alpha}\min\{\underline{D}_0, \underline{D}_1\}$. Hence it follows that

$$\begin{aligned}
d([a_0,b_0]) &= \frac{\lambda([a_0,b_i]) + \lambda([b_i, a_0+b_0-b_i]) + \lambda([a_0+b_0-b_i, b_0])}{(2(b_i-a_0)+(a_0+b_0-2b_i))^\alpha} \\
&> \frac{\lambda([a_0,b_i]) + \lambda([b_i, a_0+b_0-b_i]) + \lambda([a_0+b_0-b_i, b_0])}{2^\alpha(b_i-a_0)^\alpha+(a_0+b_0-2b_i)^\alpha} \\
&\geq \frac{\lambda([a_0,b_i]) + \lambda([b_i, a_0+b_0-b_i])}{2^\alpha(b_i-a_0)^\alpha+(a_0+b_0-2b_i)^\alpha} \\
&\geq \min\{2^{-\alpha}\frac{\lambda([a_0,b_i])}{(b_i-a_0)^\alpha}, \frac{\lambda([b_i,a_0+b_0-b_i])}{(a_0+b_0-2b_i)^\alpha}\} \\
&= \min\{2^{-\alpha}d([a_0,b_i]), d([b_i,a_0+b_0-b_i])\} \\
&\geq \min\{2^{-\alpha}\underline{D}_1, d([b_i,a_0+b_0-b_i])\}.
\end{aligned}$$

Since $[b_i, a_0+b_0-b_i] \subset [0,1]$ is an interval centered in K, the above inequality contradicts the fact that $d([a_0,b_0])$ attains the minimal value $d_{\min} < 2^{-\alpha}\underline{D}_1$. Thus the statement (1) is true.

By the statement (1) and (2), we have

$$a_0 \in [b_{i_1}, a_{i_1+1}) \text{ and } b_0 \in (b_{i_2}, a_{i_2+1}]$$

for some $1 \leq i_1 < i_2 < l_{k_0+1}$.

Since $\lambda([a_0,b_0]) = \sum_{i=i_1+1}^{i_2} \lambda([a_i,b_i])$ and $d([a_0,b_0])$ attains the minimal centered density $d_{\min}$, it follows that a_0, b_0 are taken such that (b_0-a_0) is the largest value under the condition $a_0 \in [b_{i_1}, a_{i_1+1})$, $b_0 \in (b_{i_2}, a_{i_2+1}]$ and $(a_0+b_0)/2 \in K$. Thus we have

$$b_0 - a_0 = a_{i_2+1} - b_{i_1} - 2\mathrm{dist}(\frac{a_{i_2+1}+b_{i_1}}{2}, K)$$

and

$$d([a_0,b_0]) = \frac{\sum_{i=i_1+1}^{i_2} \lambda([a_i,b_i])}{(a_{i_2+1} - b_{i_1} - 2\mathrm{dist}(\frac{a_{i_2+1}+b_{i_1}}{2}, K))^\alpha}.$$

Therefore we complete the proof of Lemma 4.21. □

Theorem 4.22.

$$d_{\min} = \min\{2^{-\alpha}\underline{D}_0, 2^{-\alpha}\underline{D}_1, D\}.$$

Here

$$\underline{D}_0 = \min\{d([0,x]) : x > 0 \text{ and } [0,x] \in \mathcal{F}_k\},$$

where k is the smallest number such that $\beta_1^{(k)} \leq \rho_1$;

$$\underline{D}_1 = \min\{d([y,1]) : y < 1 \text{ and } [y,1] \in \mathcal{F}_{k'}\},$$

where k' is the smallest number such that $\beta_{l_{k'}}^{(k')} \leq \rho_m$;

$$D = \min_{1 \leq i_1 < i_2 < l_{k_0+1}} \frac{\sum_{i=i_1+1}^{i_2} \lambda([a_i, b_i])}{(a_{i_2+1} - b_{i_1} - 2\mathrm{dist}(\frac{a_{i_2+1}+b_{i_1}}{2}, K))^\alpha},$$

where $[a_i, b_i]$ is the i-th (k_0+1)-th generation island for each $1 \leq i \leq l_{k_0+1}$.

Proof. It follows immediately from Lemma 4.11, Lemma 4.19 and Lemma 4.21. □

4.3 Examples on computing measures

In this subsection we illustrate some examples showing how to compute the Hausdorff and packing measures of linear Cantor set satisfying the generalized finite type condition with respect to the open set $(0,1)$. We will consider the examples in Section 2 again.

Example 4.23. If $\{S_j = \rho_j x + b_j\}_{j=1}^m$ satisfies the open set condition as showed in Example 2.5. We assume $0 < \rho_j < 1$ for each $j = 1, \cdots, m$. Without loss of generality, we assume that the images $S_j([0,1])$ are in increasing order, with $S_1(0) = 0$ and $S_m(1) = 1$. Then the exact α-dimensional Hausdorff measure and α-dimensional packing measure of the invariant set K can be calculated as the minimal or maximal value of a finite set of elementary functions of the parameters ρ's and b's, where α is the unique number satisfying (1.2). This is true since K naturally satisfies all the assumptions in this section. The results have already been proved in [1] and [6] for the Hausdorff measure case and packing measure case respectively. Hence our results is a naturally generalization of the results in [1, 6].

Example 4.24. Let K be the invariant set of the IFS $\{S_j\}_{j=1}^3$ on $\mathbb{R}$ defined in Example 2.6. Let α denotes the common value of the Hausdorff and packing dimensions of K. If $\rho + 2r - \rho r = 1$, then $\alpha = 1$ and $\mathcal{H}^1(K) = \mathcal{P}^1(K) = 1$. Otherwise if $\rho + 2r - \rho r < 1$, then $\alpha < 1$,

$$\underline{D}_0 = \frac{1 - r^\alpha}{(1-r)^\alpha},$$

$$\underline{D}_1 = \frac{r^\alpha}{(1 - \rho - r + \rho r)^\alpha},$$

and

$$\begin{aligned} D = \min\{ \quad & \frac{r^\alpha - r^{2\alpha}}{(\rho + r - 2\rho r - r^2 - \rho^2 + \rho^2 r - 2\mathrm{dist}(\frac{\rho + r - r^2 + \rho^2 - \rho^2 r}{2}, K))^\alpha}, \\ & \frac{r^\alpha}{(1 - r - \rho^2 - \rho r + \rho^2 r - 2\mathrm{dist}(\frac{1 - r + \rho^2 + \rho r - \rho^2 r}{2}, K))^\alpha}, \\ & \frac{2r^\alpha - r^{2\alpha}}{(1 - r^2 - \rho^2 - \rho r + \rho^2 r - 2\mathrm{dist}(\frac{1 - r^2 + \rho^2 + \rho r - \rho^2 r}{2}, K))^\alpha}, \\ & \frac{r^{2\alpha}}{(1 - r - \rho - r^2 + \rho r^2 - 2\mathrm{dist}(\frac{1 - r + \rho + r^2 - \rho r^2}{2}, K))^\alpha}, \\ & \frac{r^\alpha}{(1 - \rho - 2r^2 + \rho r^2 - 2\mathrm{dist}(\frac{1 + \rho - \rho r^2}{2}, K))^\alpha}, \\ & \frac{r^\alpha - r^{2\alpha}}{(1 - r - \rho - r^2 + \rho r - 2\mathrm{dist}(\frac{1 + r + \rho - r^2 - \rho r}{2}, K))^\alpha}\}. \end{aligned}$$

Let $k \geq 2$ be the smallest integer such that

$$2(\rho + r - \rho r) \cdot (\max\{\rho, r\})^{k-1} \leq ((1 - 2r - \rho + \rho r) \cdot \min\{\rho, r\} \cdot \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}}.$$

Then $\mathcal{H}^\alpha(K) = d_{\max}^{-1}$, where

$$d_{\max} = \max\{d(J) : J \in \mathcal{F}_k\}.$$

$\mathcal{P}^\alpha(K) = d_{\min}^{-1}$ where

$$d_{\min} = \min\{2^{-\alpha}\underline{D}_0, 2^{-\alpha}\underline{D}_1, D\}.$$

Proof. If $\rho + 2r - \rho r = 1$, then $K = [0, 1]$, which yields the trivial result. Hence we only need to consider the second case, i.e., $\rho + 2r - \rho r < 1$. In this case, k_0=1. Using Lemma 4.11, we get

$$\underline{D}_0 = \min\{d([0, x]) : x > 0 \text{ and } [0, x] \in \mathcal{F}_2\},$$

and

$$\underline{D}_1 = \min\{d([y, 1]) : y < 1 \text{ and } [y, 1] \in \mathcal{F}_1\}.$$

We adopt the same setup and notations of Example 2.6, with $\mathcal{T}_1 = [I_0]$ and $\mathcal{T}_2 = [I_1]$ where $I_0 = [0, 1]$ is the root island and $I_1 = S_1([0, 1]) \cup S_2([0, 1])$. Using the definition of λ, we have $\lambda(I_0) = 1$ and $\lambda(I_1) = 1 - r^\alpha$. Then after a detailed computation, we get the exact values

$$\underline{D}_0 = \min\{1, \frac{1 - r^\alpha}{(1 - r)^\alpha}, \frac{(\rho^\alpha + r^\alpha)(1 - r^\alpha)}{(\rho + r)^\alpha (1 - r)^\alpha}, \frac{1 - r^{2\alpha}}{(1 - r^2)^\alpha}\} = \frac{1 - r^\alpha}{(1 - r)^\alpha},$$

and

$$\underline{D}_1 = \min\{1, \frac{r^\alpha}{(1-\rho-r+\rho r)^\alpha}\} = \frac{r^\alpha}{(1-\rho-r+\rho r)^\alpha}.$$

Hausdorff measure. A detailed calculation shows

$$\gamma_{\min}^{(k_0+1)} = \gamma_{\min}^{(2)} = (1-2r-\rho+\rho r)\cdot \min\{\rho, r\},$$

and

$$\beta_{\max}^{(k)} = (\rho + r - \rho r)\cdot (\max\{\rho, r\})^{k-1}$$

for each $k \geq 1$. Let $k \geq 2$ be the smallest integer such that

$$2\beta_{\max}^{(k)} \leq (\gamma_{\min}^{(2)} \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}}.$$

Then by Theorem 4.14, the maximal density $d_{\max}$ is attained for an interval in $\mathcal{F}_k$. Furthermore, by Corollary 1.3, $\mathcal{H}^\alpha(K) = d_{\max}^{-1}$.

Packing measure. Since $k_0 = 1$, the constant D in Theorem 4.22 is

$$D = \min_{1\leq i_1 < i_2 < 5} \frac{\sum_{i=i_1+1}^{i_2} \lambda([a_i, b_i])}{(a_{i_2+1} - b_{i_1} - 2\mathrm{dist}(\frac{a_{i_2+1}+b_{i_1}}{2}, K))^\alpha}.$$

where $[a_i, b_i]$ is the i-th 2-th generation island for each $1 \leq i \leq 5$. A detailed calculation yields the exact value of D. Then by Theorem 4.22, the minimal centered density

$$d_{\min} = \min\{2^{-\alpha}\underline{D}_0, 2^{-\alpha}\underline{D}_1, D\}.$$

Hence by Corollary 1.4, $\mathcal{P}^\alpha(K) = d_{\min}^{-1}$. □

Remark 4.25. Consider the special case where $\rho = r = 1/16$ in the above example. By Example 2.13, the dimension $\alpha = \log_{16} 2/(3-\sqrt{5}) \approx 0.3471$. And we calculate that $\underline{D}_0 = (16^\alpha - 1)/15^\alpha \approx 0.6320$, $\underline{D}_1 = 16^\alpha/225^\alpha \approx 0.3995$, and $\gamma_{\min}^{(2)} = 209/4096 \approx 0.0510$. Moreover, for each $k \geq 1$, $\beta_{\max}^{(k)} = (31/256)\cdot(1/16^{k-1})$. Hence the smallest k should satisfy

$$2\cdot\frac{31}{256}\cdot\frac{1}{16^{k-1}} \leq (0.0510\cdot 0.3995)^{\frac{1}{1-\alpha}},$$

which yields that $k = 3$. After a complicated computation by computer, we eventually get $d_{\max} = (256^\alpha - 16^\alpha)/31^\alpha \approx 1.2861$, and $\mathcal{H}^\alpha(K) = 31^\alpha/(256^\alpha - 16^\alpha) \approx 0.7775$.

For the packing measure, a detailed calculation by computer shows that $D = 8^\alpha/225^\alpha \approx 0.3140$. Hence $d_{\min} = 2^{-\alpha}\underline{D}_1 = D = 8^\alpha/225^\alpha \approx 0.3140$ and $\mathcal{P}^\alpha(K) = 225^\alpha/8^\alpha \approx 3.1843$.

Example 4.26. Let K be the invariant set of the IFS $\{S_j\}_{j=1}^3$ on $\mathbb{R}$ defined in Example 2.8. Let α denotes the common value of the Hausdorff and packing dimensions of K. Then α is the logarithmic ratio of the largest root of the polynomial equation

$$x^3 - 6x^2 + 5x - 1 = 0$$

to 9, $\alpha \approx 0.7369$. Moreover,

$$\mathcal{H}^\alpha(K) = \frac{11^\alpha}{27^\alpha - 9^\alpha} \approx 0.9297,$$

and

$$\mathcal{P}^\alpha(K) = \frac{32^\alpha}{9^\alpha} \approx 2.5467.$$

Proof. The dimension result is immediately get from Example 2.14. We adopt the same setup and notations of Example 2.8, with $\mathcal{T}_1 = [I_0]$, $\mathcal{T}_2 = [I_1]$ and $\mathcal{T}_3 = [I_2]$ where $I_0 = [0,1]$ is the root island, $I_1 = S_{11}([0,1]) \cup S_{12}([0,1])$ and $I_2 = S_{13}([0,1]) \cup S_2([0,1])$. Using the definition of λ, we have $\lambda(I_0) = 1$, $\lambda(I_1) = 1/3^\alpha - 1/9^\alpha$ and $\lambda(I_2) = 1 - 2/3^\alpha + 1/9^\alpha$. Using Lemma 4.11, we get

$$\underline{D}_0 = \min\{d([0,x]) : x > 0 \text{ and } [0,x] \in \mathcal{F}_1\},$$

and

$$\underline{D}_1 = \min\{d([y,1]) : y < 1 \text{ and } [y,1] \in \mathcal{F}_1\}.$$

A detailed calculation shows that

$$\underline{D}_0 = \min\{1, \frac{3^\alpha - 1}{2^\alpha}, \frac{9^\alpha - 1}{8^\alpha}\} = \frac{3^\alpha - 1}{2^\alpha} \approx 0.7482,$$

and

$$\underline{D}_1 = \min\{1, \frac{9^\alpha}{16^\alpha}, \frac{81^\alpha - 27^\alpha + 9^\alpha}{70^\alpha}\} = \frac{9^\alpha}{16^\alpha} \approx 0.6544.$$

Hausdorff measure. Since $k_0 = 1$, a detailed calculation shows

$$\gamma_{\min}^{(k_0+1)} = \gamma_{\min}^{(2)} = \frac{7}{729} \approx 0.0096.$$

By Theorem 4.14, we need to find a smallest integer k such that

$$2\beta_{\max}^{(k)} \leq (\gamma_{\min}^{(2)} \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}} = (\frac{7}{729} \min\{\frac{3^\alpha - 1}{2^\alpha}, \frac{9^\alpha}{16^\alpha}\})^{\frac{1}{1-\alpha}}.$$

Noticing that $\beta_{\max}^{(k)} = 5/(27 \cdot 9^{k-1})$, the smallest $k = 10$.

Then by Theorem 4.12, the maximal density $d_{\max}$ is attained for an interval in $\mathcal{F}_{10}$. Furthermore, by Corollary 1.3, $\mathcal{H}^\alpha(K) = d_{\max}^{-1}$. After a complicated computation by computer, we eventually get $d_{\max} = (27^\alpha - 9^\alpha)/11^\alpha \approx 1.0756$, and $\mathcal{H}^\alpha(K) = 11^\alpha/(27^\alpha - 9^\alpha) \approx 0.9297$.

Packing measure. Since $k_0 = 1$, the constant D in Theorem 4.22 is

$$D = \min_{1 \leq i_1 < i_2 < 20} \frac{\sum_{i=i_1+1}^{i_2} \lambda([a_i, b_i])}{(a_{i_2+1} - b_{i_1} - 2\mathrm{dist}(\frac{a_{i_2+1} + b_{i_1}}{2}, K))^\alpha}.$$

where $[a_i, b_i]$ is the i-th 2-th generation island for each $1 \le i \le 20$. A detailed calculation by computer yields the exact value of $D = 9^\alpha/32^\alpha \approx 0.3927$. Hence $d_{\min} = 2^{-\alpha}\underline{D}_1 = D = 9^\alpha/32^\alpha \approx 0.3927$ and $\mathcal{P}^\alpha(K) = 32^\alpha/9^\alpha \approx 2.5467$. □

Although these algorithms applies in theory to any case considered under Assumption A and Assumption B, in practice it is useable in very few cases. Even in the following simple example.

Example 4.27. Let K be the invariant set of the IFS $\{S_j\}_{j=1}^4$ on $\mathbb{R}$ defined in Example 2.9. Let α denotes the common value of the Hausdorff and packing dimensions of K. Then

$$\alpha = \log_4(5+\sqrt{5}) - \frac{1}{2} \approx 0.9276,$$

and

$$\mathcal{P}^\alpha(K) = 3^\alpha \approx 2.7706.$$

However, our algorithm for computing $\mathcal{H}^\alpha(K)$ exceeds the computing power.

Proof. The dimension result is immediately get from Example 2.15. We adopt the same setup and notations of Example 2.9, with $\mathcal{T}_1 = [I_0]$, $\mathcal{T}_2 = [I_2]$ and $\mathcal{T}_3 = [I_5]$ where $I_0 = [0,1]$ is the root island, $I_2 = S_2([0,1]) \cup S_3([0,1])$ and $I_5 = S_{22}([0,1]) \cup S_{23}([0,1]) \cup S_{31}([0,1])$. Using the definition of λ, we have $\lambda(I_0) = 1$, $\lambda(I_2) = 1 - 2/4^\alpha$ and $\lambda(I_5) = 1 - 3/4^\alpha$. Using Lemma 4.11, we get

$$\underline{D}_0 = \min\{d([0,x]) : x > 0 \text{ and } [0,x] \in \mathcal{F}_1\},$$

and

$$\underline{D}_1 = \min\{d([y,1]) : y < 1 \text{ and } [y,1] \in \mathcal{F}_1\}.$$

A detailed calculation shows that

$$\underline{D}_0 = \min\{1, \frac{4^\alpha - 1}{3^\alpha}\} = \frac{4^\alpha - 1}{3^\alpha} \approx 0.9449,$$

and

$$\underline{D}_1 = \min\{1, \frac{2^\alpha}{3^\alpha}, \frac{4^\alpha - 1}{3^\alpha}\} = \frac{2^\alpha}{3^\alpha} \approx 0.6865.$$

Hausdorff measure. We will use Theorem 4.15 since there exist touching islands and $\rho_1 = \rho_m$. Observe that $k_0 = 2$, $n_1 = n_m = 1$ and $\eta = 1/8$. A detailed calculation shows

$$\beta_{\min}^{(k_0+1)} = \beta_{\min}^{(3)} = \frac{1}{64}.$$

By Theorem 4.15, we need to find a smallest integer k such that

$$\begin{aligned} 2\beta_{\max}^{(k)} &\le (\eta \beta_{\min}^{(k_0+1)} \rho_1^{n_1} \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}} \\ &= (\frac{1}{8} \cdot \frac{1}{64} \cdot \frac{1}{4} \min\{\frac{4^\alpha - 1}{3^\alpha}, \frac{2^\alpha}{3^\alpha}\})^{\frac{1}{1-\alpha}}. \end{aligned}$$

Noticing that $\beta_{\max}^{(k)} = 1/(8 \cdot 4^{k-2}) = 1/2^{2k-1}$, the smallest $k = 81$.

Then by Theorem 4.15, the maximal density $d_{\max}$ is attained for an interval in $\mathcal{F}_{81}$. Furthermore, by Corollary 1.3, $\mathcal{H}^\alpha(K) = d_{\max}^{-1}$. However, the time involved in searching all sets in $\mathcal{F}_k$ rapidly becomes impractical.

Packing measure. Since $k_0 = 2$, the constant D in Theorem 4.22 is

$$D = \min_{1 \le i_1 < i_2 < 35} \frac{\sum_{i=i_1+1}^{i_2} \lambda([a_i, b_i])}{(a_{i_2+1} - b_{i_1} - 2\mathrm{dist}(\frac{a_{i_2+1}+b_{i_1}}{2}, K))^\alpha}.$$

where $[a_i, b_i]$ is the i-th 3-th generation island for each $1 \le i \le 35$. A detailed calculation by computer yields the exact value of $D = 1/3^\alpha \approx 0.3609$. Hence $d_{\min} = 2^{-\alpha}\underline{D}_1 = D = 1/3^\alpha \approx 0.3609$ and $\mathcal{P}^\alpha(K) = 3^\alpha \approx 2.7706$. □

5 Further Discussions

Are Assumption A and Assumption B necessary?

If we permit the IFS not to satisfy Assumption A or Assumption B, things become more complicated. It seems hard to get uniform formulae for $\mathcal{H}^\alpha(K)$ and $\mathcal{P}^\alpha(K)$. However, in some special cases, we can still use the similar method to get the similar measure results. The following are two concrete examples.

Example 5.1. Consider the IFS $\{S_j\}_{j=1}^3$ as follows.

$$S_1(x) = \frac{1}{3}x, \quad S_2(x) = \frac{1}{9}x + \frac{2}{9}, \quad S_3(x) = \frac{1}{3}x + \frac{2}{3}.$$

If we choose $\mathcal{M}_k = \Lambda_k$ for each $k \ge 0$, $\{S_j\}_{j=1}^3$ is of generalized finite type with respect to the invariant set $(0,1)$, with Assumption A not satisfied. See Figure 5.

Since the proof is obvious, we omit it. In fact, the invariant set K of this IFS is the classical Cantor ternary set. It is well-known that the common value of the Hausdorff and packing dimensions of K is equal to $\log_3 2$ and the $\log_3 2$-dimensional Hausdorff measure of K is equal to 1 (see [1]). Replacing the IFS by $\{S_1, S_3\}$ equivalently, from Theorem 4.22, one gets that $\mathcal{P}^{\log_3 2} = 4^{\log_3 2}$. (See this also in [6].)

Example 5.2. Consider the IFS $\{S_j\}_{j=1}^3$ defined as follows.

$$S_1(x) = \rho x, \quad S_2(x) = \rho x + \rho\frac{1-\rho}{1+\rho}, \quad S_3(x) = \rho x + 1 - \rho,$$

where $0 < \rho < 1/3$. Choose $\mathcal{M}_k = \Sigma_k$ for each $k \ge 0$, then $\{S_j\}_{j=1}^3$ is of generalized finite type with respect to the invariant set $(0,1)$ with Assumption B not satisfied. The common



(a) (b)

Figure 5. (a) The first five levels of islands in Example 5.1. Note that vertical bars have been added at island boundaries. (b) How the distinct overlap types are generated in Example 5.1. Iterates of the point 0 *under the similitudes are represented by dots.*

value of the Hausdorff and packing dimensions of the invariant set K is $\alpha = \log_\rho 2/(3+\sqrt{5})$. Moreover, let $k \geq 2$ be the smallest integer such that

$$4\rho^k \leq (\frac{(1-3\rho)\rho^{1+\alpha}}{(1-\rho)^\alpha})^{\frac{1}{1-\alpha}}.$$

Then $\mathcal{H}^\alpha(K) = d_{\max}^{-1}$, where

$$d_{\max} = \max\{d(J) : J \in \mathcal{F}_k\}.$$

$\mathcal{P}^\alpha(K) = d_{\min}^{-1}$ where

$$d_{\min} = \min\{\frac{\rho^\alpha(1+\rho)^\alpha}{2^\alpha(1-\rho)^\alpha}, D\},$$

and

$$\begin{aligned} D = \min\{ \quad & \frac{\rho^\alpha - \rho^{2\alpha}}{(\frac{2\rho-3\rho^2-\rho^3}{1+\rho} - 2\mathrm{dist}(\frac{2\rho+\rho^2-\rho^3}{2(1+\rho)}, K))^\alpha}, \frac{\rho^\alpha}{(\frac{1-3\rho^2}{1+\rho} - 2\mathrm{dist}(\frac{1+\rho^2}{2(1+\rho)}, K))^\alpha}, \\ & \frac{2\rho^\alpha - \rho^{2\alpha}}{(\frac{1+\rho-3\rho^2-\rho^3}{1+\rho} - 2\mathrm{dist}(\frac{1+\rho+\rho^2-\rho^3}{2(1+\rho)}, K))^\alpha}, \frac{\rho^{2\alpha}}{(1-2\rho-2\mathrm{dist}(\frac{1}{2}, K))^\alpha}, \\ & \frac{\rho^\alpha}{(1-\rho-\rho^2-2\mathrm{dist}(\frac{1+\rho-\rho^2}{2}, K))^\alpha}, \frac{\rho^\alpha - \rho^{2\alpha}}{(\frac{1-\rho-\rho^2-\rho^3}{1+\rho} - 2\mathrm{dist}(\frac{1+3\rho-\rho^2-\rho^3}{2(1+\rho)}, K))^\alpha}\}. \end{aligned}$$

Proof. The proof of that $\{S_j\}_{j=1}^3$ is of generalized finite type with respect to the invariant set $(0,1)$ is similar to the proof of Example 2.6, in which S_{21}, S_{22} are replaced by



(a) (b)

Figure 6. (a) The first five levels of islands in Example 5.2 with parameters $\rho = 1/6$*. Note that vertical bars have been added at island boundaries. (b) How the distinct overlap types are generated in Example 5.2. Iterates of the point* 0 *under the similitudes are represented by dots.*

S_{22}, S_{21} respectively. Observing that $\rho(1-\rho)/(1+\rho) \in S_1([0,1]) \cap K$ and $\rho(1-\rho)/(1+\rho) \notin S_1K$, one gets that Assumption B is not satisfied. See Figure 6.

The dimension result is immediately get by Theorem 2.10. In the following, we turn to prove the measure results.

It is easy to verify that $[I_0]$ and $[I_1]$ denoted respectively by $\mathcal{T}_1$ and $\mathcal{T}_2$ are the all distinct overlap types, where $I_0 = [0,1]$ and $I_1 = S_1([0,1]) \cup S_2([0,1])$. It is worth mentioning that in this case Lemma 4.10 may not hold since Assumption B is not satisfied. Hence the original notations $\underline{D}_0$ and $\underline{D}_1$ may not suitable. However, we redefine them by $\underline{D}_0 := \min\{\underline{D}_0^1, \underline{D}_0^2\}$, and $\underline{D}_1 := \min\{\underline{D}_1^1, \underline{D}_1^2\}$. A similar discussion as the proof of Lemma 4.10 shows that $\underline{D}_0^1 \le \underline{D}_0^2$ and $\underline{D}_1^1 \le \underline{D}_1^2$. Moreover, observing the distribution of the offsprings of I_1, we can also get $\underline{D}_1^2 \le \underline{D}_1^1$. Hence $\underline{D}_0 = \underline{D}_0^1$ and $\underline{D}_1 = \underline{D}_1^1 = \underline{D}_1^2$. By a similar argument of Lemma 4.11 (with suitable modifications), we get

$$\underline{D}_0 = \min\{d([0,x]) : x > 0 \text{ and } [0,x] \in \mathcal{F}_2\},$$

and

$$\underline{D}_1 = \min\{d([y,1]) : y < 1 \text{ and } [y,1] \in \mathcal{F}_1\}.$$

Using the definition of λ, we have $\lambda(I_0) = 1$ and $\lambda(I_1) = 1 - \rho^\alpha$. Hence we can get that $d([0,x])$ attains the minimal value

$$\underline{D}_0 = \min\{\frac{2(1-\rho^\alpha)(1+\rho)^\alpha}{(1-\rho)^\alpha(2+\rho)^\alpha}, \frac{1-\rho^\alpha}{(1-\rho)^\alpha}\} = \frac{1-\rho^\alpha}{(1-\rho)^\alpha}$$

at the point $x_0 = 1 - \rho$, and $d([y, 1])$ attains the minimal value

$$\underline{D}_1 = \rho^\alpha \frac{(1+\rho)^\alpha}{(1-\rho)^\alpha}$$

at the point $y_0 = 2\rho/(1+\rho)$.

Hausdorff measure. $k_0 = 1$, $\gamma_{\min}^{(2)} = \rho(1-3\rho)/(1+\rho)$, $\beta_{\max}^{(k)} = 2\rho^k/(1+\rho)$ for each $k \geq 1$. By a suitable modification of Theorem 4.14, we get $d_{\max}$ is attained for an interval in $\mathcal{F}_k$ where $k \geq 2$ is the smallest integer such that

$$2\beta_{\max}^{(k)} \leq (\gamma_{\min}^{(2)} \min\{\underline{D}_0, \underline{D}_1\})^{\frac{1}{1-\alpha}}.$$

Furthermore, by Corollary 1.3, $\mathcal{H}^\alpha(K) = d_{\max}^{-1}$.

Packing measure. We need a similar result of Lemma 4.19, i.e., $d_{\min} \leq 2^{-\alpha} \min\{\underline{D}_0, \underline{D}_1\}$. However, at first glance, we can not prove it in general. The reason is the following. Recall the proof of Lemma 4.19, we should find two intervals J_0 and J_1 centered in K with $d(J_0) = 2^{-\alpha}\underline{D}_0$ and $d(J_1) = 2^{-\alpha}\underline{D}_1$. In fact, we can define J_1 as that used in the proof of Lemma 4.19 since $\underline{D}_1^1 = \underline{D}_1^2$ and $S_3([0,1]) \cap K = S_3K$. (Half of Assumption B holds.) But for J_0, the original process is invalid since $\underline{D}_0^1$ and $\underline{D}_0^2$ may be not equal. Fortunately, a detailed verifying shows that the inequality $\underline{D}_1 \leq \underline{D}_0$ always holds, which ensures that it is meaningless to find J_0, i.e., the existence of J_1 is enough. Hence Lemma 4.19 remains true. Once Lemma 4.19 is proved, Lemma 4.20, Lemma 4.21 and eventually Theorem 4.22 are followed automatically, in which $\min\{\underline{D}_0, \underline{D}_1\}$ are all replaced by $\underline{D}_1$. $\square$

Remark 5.3. Consider the special case where $\rho = 1/6$ in the above example. The dimension $\alpha = \log_6((3+\sqrt{5})/2) \approx 0.5371$. And we calculate that $\underline{D}_0 = (6^\alpha - 1)/5^\alpha \approx 0.6816$, $\underline{D}_1 = 7^\alpha/30^\alpha \approx 0.4576$, and $\gamma_{\min}^{(2)} = 1/14 \approx 0.0714$. Moreover, for each $k \geq 1$, $\beta_{\max}^{(k)} = (2/7) \cdot (1/6^{k-1})$. Hence the smallest k should satisfy

$$2 \cdot \frac{2}{7} \cdot \frac{1}{6^{k-1}} \leq (0.0714 \cdot 0.4576)^{\frac{1}{1-\alpha}},$$

which yields that $k = 5$. After a complicated computation by computer, we eventually get $d_{\max} = 2(6^\alpha - 1)/6^\alpha \approx 1.2361$, and $\mathcal{H}^\alpha(K) = 6^\alpha/(2(6^\alpha - 1)) \approx 0.8090$.

For the packing measure, a detailed calculation by computer shows that $D = 7^\alpha/60^\alpha \approx 0.3154$. Hence $d_{\min} = 2^{-\alpha}\underline{D}_1 = D = 7^\alpha/60^\alpha \approx 0.3154$ and $\mathcal{P}^\alpha(K) = 60^\alpha/7^\alpha \approx 3.1709$.

Can we allow negative ρ's?

Consider the IFSs containing orientation reversing similarities, which simply means that we allow some of the ρ_j to be negative. In some special cases, it can be proved by using a similar method that the results of Section 4 still remain true.

We again normalize by assuming that $S_j([0,1])$ are in increasing order with $S_1([0,1])$ containing 0 and $S_m([0,1])$ containing 1. For example, $\rho_1\rho_m > 0$, $\bigcup_{2\leq j\leq m} S_j([0,1]) \cap$

$S_1([0,1]) = \emptyset$, $\bigcup_{1\le j\le m-1} S_j([0,1]) \cap S_m([0,1]) = \emptyset$, and $\gamma_{\min}^{(k_0+1)} > 0$. Obviously, in this case Assumption B is naturally satisfied. We can still obtain measure results following the idea of Section 4.

To illustrate this, we assume that ρ_1 and ρ_m are both positive since otherwise we can replace the IFS with its iterated square, i.e., all compositions S_iS_j. Similar to Example 5.2, for this case, Lemma 4.10 may not hold. Hence the original notations $\underline{D}_0$ and $\underline{D}_1$ will be meaningless. In spite of this, the following constant defined by $\underline{D} = \min\{\underline{D}_0^1, \underline{D}_1^1\}$ will replace the important role of the original notation $\min\{\underline{D}_0, \underline{D}_1\}$ in Section 4. It is not hard to verify that $\underline{D} \le \underline{D}_0^i$ and $\underline{D} \le \underline{D}_1^i$ for all $1 \le i \le q$, where q is the cardinality of all the distinct overlap types. Moreover, we could also characterize the constant $\underline{D}$ using a suitable modification of Lemma 4.11, i.e.,

$$\underline{D} = \min\{d(J) : J \in \mathcal{F}_1 \text{ and } J \text{ is of the form } [0,x] \text{ or } [y,1]\}.$$

It is not hard to verify that Theorem 4.14 for Hausdorff measure, and Lemma 4.19, Lemma 4.20, Lemma 4.21 and Theorem 4.22 for packing measure still remain true in which $\min\{\underline{D}_0, \underline{D}_1\}$ is replaced by $\underline{D}$.

How about self-similar sets in higher dimensional Euclidean spaces?

As mentioned earlier, with suitable modifications if necessary, the results in Section 3 may be generalized to self-similar sets in higher dimensional Euclidean spaces. How about measure results in Section 4? Our answer is: almost all the obvious generalizations of our results are false.

It is clear that the blow-up principles continue to hold. So we can also focus attention to sets not contained entirely in some (k_0+1)-generation island, where k_0 is also the smallest non-negative integer such that none of the islands in $\mathcal{F}_{k_0+1}$ is of a new overlap type. For the maximal density $d_{\max}$, since any set can be replaced by its convex hull without decreasing the density, it is reasonable to limit any searching algorithm to convex sets. However, it does not follow that the maximal density among $\mathcal{F}_k$ sets is achieved by a convex set, since the convex hull of a set in $\mathcal{F}_k$ may not belong to $\mathcal{F}_k$. To illustrate this, in [1] Ayer & Strichartz consider a concrete example, i.e, the usual Sierpinski gasket in plane. We omit it here. For the minimal centered density $d_{\min}$, one obvious obstacle is that almost all lemmas concerning $d_{\min}$ require $\alpha < 1$. Moreover, even if we were to limit attention to self-similar sets of dimension $\alpha < 1$, it is unlikely that the same results would hold. In fact, we would have to confront how to describe $\underline{D}_0$ and $\underline{D}_1$ and how to overcome the difficulty of the calculation of densities of higher dimensional sets. (It seems uncontrollable.)

References

[1] E. Ayer, R. Strichartz, Exact Hausdorff measure and intervals of maximum density for Cantor sets, Trans. Amer. Math. Soc. 351 (1999) 3725-3741.

[2] K.J. Falconer, The Geometry of Fractal Sets, Cambridge Tracts in Mathematics 85, Cambridge Univ. Press, New York-London, 1985.

[3] K.J. Falconer, Dimensions and measures of quasi self-similar sets, Proc. Amer. Math. Soc. 106 (1989) 543-554.

[4] K.J. Falconer, Fractal Geometry. Mathematical Foundations and Applications, John Wiley & Sons, Ltd., Chichester, 1990.

[5] D.-J. Feng, S. Hua and Z.-Y. Wen. Some relations between packing premeasure and packing measure, Bull. London Math. Soc. 31 (1999) 665-670.

[6] D.-J. Feng, Exact packing measure of Cantor sets, Math. Natchr. 248-249 (2003) 102-109.

[7] J.E. Hutchinson, Fractals and self-similarity, Indiana Univ. Math. J. 30 (1981) 713-747.

[8] S.P. Lalley, β-expansions with deleted digits for Pisot numbers, Trans. Amer. Math. Soc. 349 (1997) 4355-4365.

[9] K.-S. Lau, S.-M. Ngai, A generalized finite type condition for iterated function systems, Adv. Math. 208 (2007) 647-671.

[10] K.-S. Lau, X.-Y. Wang, Iterated function systems with a weak separation condition, Studia Math. 161 (3) (2004) 249-268.

[11] J. Marion, Measure de Hausdorff d'un fractal à similitude interne, Ann. Sci. Math. Québec 10 (1986) 51-84.

[12] J. Marion, Measures de Hausdorff d'ensembles fractals, Ann. Sci. Math. Québec 11 (1987) 111-132.

[13] P. Mattila, Geometry of Sets and Measures in Euclidean Spaces, in: Fractals and Rectifiability, Cambridge University Press, 1995.

[14] R.D. Mauldin, S.C. Williams, Hausdorff dimension in graph directed constructions, Trans. Amer. Math. Soc. 309 (1988) 811-829.

[15] R.D. Mauldin, M. Urbański, Dimensions and measures in infinite iterated function systems, Proc. London Math. Soc. (3)73 (1996) 105-154.

[16] P.A.P. Moran, Additive functions of intervals and Hausdorff measure, Proc. Cambridge Philos. Soc. 42 (1946) 15-23.

[17] M. Morán, Computability of the Hausdorff and pakcing measures on self-similar sets and the self-similar tiling principle, Nonlinearity 18 (2005) 559-570.

[18] S.-M. Ngai, Y. Wang, Hausdorff dimension of self-similar sets with overlaps, J. London Math. Soc. (2)63 (2001) 655-672.

[19] L. Olsen, Density theorems for Hausdorff and packing measures of self-similar sets, Aequationes Math. 75 (2008) 208-225.

[20] H. Qiu, Continuity of packing measure function of self-similar iterated function systems, to appear in Ergod. Th. & Dynam. Sys.

[21] H. Rao, Z.-Y. Wen, A class of self-similar fractals with overlap structure, Adv. in Appl. Math. 20 (1998) 50-72.

[22] A. Schief, Separation properties for self-similar sets, Proc. Amer. Math. Soc. 122 (1994), 111-115.

[23] S.J. Taylor, C. Tricot, Packing measure, and its evaluation for a Brownian path, Trans. Amer. Math. Soc. 288 (1985) 679-699.

[24] C. Tricot, Two definitions of fractional dimension, Math. Proc. Camb. Phil. Soc. 91 (1982), 54-74.

[25] Z.-L. Zhou, L. Feng, Twelve open questions on the exact value of the Hausdorff measure and on the topogical entropy: a brief review of recent results, Nonlinearly. 17 (2004) 493-502.